

Strategy-Proof Exchange under Trichotomous Preferences

Vikram Manjunath* Alexander Westkamp†

November 28, 2018

Abstract

We study the exchange of indivisible objects without monetary transfers when each agent may be endowed with and consume more than one object. We assume that each agent has *trichotomous* preferences in the sense that she

- (A) partitions objects into desirable and undesirable ones,
- (B) considers a bundle unacceptable if it contains undesirable objects that she was not already endowed with, and
- (C) ranks acceptable bundles by their numbers of desirable objects.

On this domain, we show that there is an individually rational, Pareto-efficient, and strategy-proof mechanism that is also computationally efficient.

JEL classification: C78, D82

Keywords: individual rationality, Pareto-efficiency, strategy-proofness, indivisible goods, multi-unit demand

1 Introduction

We study the problem of reallocating indivisible objects without monetary transfers. Unlike much of the earlier work on this problem,¹ we consider situations where each agent may be

*Department of Economics, University of Ottawa; e-mail: vikram@dosamobile.com.

†Department of Management, Economics and Social Sciences, University of Cologne; e-mail: westkamp@wiso.uni-koeln.de.

¹See the literature following [Shapley and Scarf \[1974\]](#).

endowed with and consume more than one object. We require exchange to be balanced in the sense that each agent ends up with exactly the same number of objects as she is endowed with. Agents' preferences over individual objects are coarse: An object is either desirable or undesirable. Each agent considers a bundle unacceptable if it contains an undesirable objects that she was not endowed with. Finally, each agent ranks acceptable bundles in increasing order of the numbers of desirable objects that they contain. Our contribution is to define an individually rational, Pareto-efficient, and strategy-proof mechanism that is computationally efficient. Our positive result is in marked contrast with the impossibility results that abound in the literature on multi-object exchange without monetary transfers [Sönmez, 1999, Biró et al., 2018].

Our interest in these exchange problems stems from our search for a solution to the problem of shift-reallocation. Millions of people in many different professions, from physicians to retail workers, engage in shift work. Shift plans are often made months in advance and scenarios like the following are common: A medical practice consists of four specialist consultants Drs A, B, C, and D. This practice is responsible for ensuring that emergency medical services in their specialty are available to a given hospital at all times. That is, each week, one of the four doctors is to be designated as being *on call*. Being on call is not desirable for these doctors. However, it is necessary for their practice to maintain privileges at the hospital. Since it is a chore that they must perform, in the interest of fairness, they agree to share the weeks equally so that each member of the group is on call every fourth week.² To facilitate planning, the call schedule is made six months at a time, taking the doctors' preferences into consideration. For instance, the schedule from January 1 to June 30 is announced in December, and is created on the basis of the doctors' preferences as of December. Thus, on January 1, each of the four doctors is responsible for their assigned weeks until June 30. While this initial assignment may be Pareto-efficient with regards to the doctors' December preferences, six months is a long time. At some later point, say the beginning of March, there may be scope for re-optimization based on current preferences. It may well happen that Dr. A would like to attend a conference the week of April 5, Dr. D would like to help at a clinic in a remote area on June 16, and Dr. C would like to go on vacation on May 23. If each of these doctors is obliged to be on call for the respective week, a three way trade could improve welfare in regards to current (as of March) preferences. We are interested in the design of mechanisms to identify such trades optimally and provide agents with incentives to reveal their private information about scheduling conflicts truthfully.³

²In the case of medical residents, such a restriction would not only be for reasons of fairness, but also for training purposes.

³Note that the re-optimization at the beginning of March in the above example is a static problem. We do not consider the issue of optimally timed matching as in Akbarpour et al. [2018].

The novel aspect of the model that we study is in agents’ preferences. First, each agent partitions the set of all objects into desirable and undesirable ones. Second, no agent is ever willing to accept, i.e. would rather stick to her endowment, a bundle that contains one or more undesirable objects that she is not endowed with. Finally, each agent ranks acceptable bundles in increasing order of the numbers of desirable objects they contain. We call such preferences *trichotomous*. In the context of shift exchange, a shift is undesirable (desirable) to a worker if it is (not) in conflict with any other activities (e.g. paid work for other firms, further training/education, vacation) that the worker would like to or has already committed to. Furthermore, a worker is never willing to participate in an exchange that creates new scheduling conflicts in the sense that she is assigned new undesirable shifts. Put differently, workers will always find a way to fulfil the obligations associated with the initial shift schedule, while they face some hard constraints that limit which alternative shifts they can take on.

We focus on situations in which endowments are commonly known and exchange is balanced in the sense that each agent ends up with the same number of goods as she was endowed with. In the context of shift exchange, this means that all participants know the initial schedule of shifts and that a worker’s total workload is fixed at the number of shifts in the initial schedule. Given that preferences are trichotomous, the crucial piece of information that we need to elicit from an agent is which objects she finds desirable. An exchange mechanism asks each agent to reveal her set of desirable objects and then computes the final allocation of shifts on the basis of agents’ reports as well as their commonly known endowments. We are interested in mechanisms that satisfy the following three desiderata: *individual rationality* (no agent is assigned an unacceptable bundle), *Pareto-efficiency* (no further reallocation can make any agent better off without harming another), and *strategy-proofness* (no agent can ever profit from lying about her set of desirable objects). The main contribution of this paper is to define a new class of balanced exchange mechanisms satisfying all of these desiderata.⁴ Our *Individually Rational Priority* (IRP) mechanisms are parameterized by a fixed ordering of the agents, independent of their reports, and pick a final allocation of goods in accordance with the following sequential process:

- In the first step, find the maximum number of desirable objects that the first agent can receive in an individually rational allocation. Call this number the “promise” to the first agent.
- In the t^{th} step, find the maximum number of desirable objects that the t^{th} agent can receive in an individually rational allocation subject to the constraints imposed by the

⁴For mechanisms that solve the unit endowment problem under general weak preferences, see [Jaramillo and Manjunath \[2012\]](#), [Alcalde-Unzu and Molis \[2011\]](#), [Aziz and De Keijzer \[2012\]](#), and [Saban and Sethuraman \[2013\]](#).

promises made to the first $t - 1$ agents. Call this number the “promise” to the t^{th} agent.

An IRP mechanism selects an outcome that meets the promises made to all of the agents in this process. We show that, given an ordering, an IRP mechanism is individually rational, Pareto-efficient, and strategy-proof. While individual rationality and Pareto-efficiency are almost by definition, it is significantly harder to show that this mechanism makes truthful revelation a weakly dominant strategy for the agents. The key feature of an IRP mechanism that makes establishing strategy-proofness difficult is that the report of an agent alters the set of individually rational allocations and can thereby affect the outcomes for all agents.⁵ An agent who is not first in line may therefore hope that she can influence the promises to her predecessors in such a way that the mechanism promises her more truly desirable objects. In our proof, we develop an elaborate combinatorial argument to show that such hopes are misguided.

Apart from their desirable allocative and incentive properties, we also show that IRP mechanisms are computationally efficient. More precisely, taking agents’ desirable sets and a priority order over the agents as inputs, an IRP allocation may be computed in $O(mn^3)$ time, where m is the total number of objects and n is the number of agents.

Related Literature

Without rather strong restrictions on preferences, individual rationality, Pareto-efficiency, and strategy-proofness are incompatible when agents are endowed with and demand multiple objects [Sönmez, 1999, Biró et al., 2018]. In a companion paper [Manjunath and Westkamp, 2018], we show that even for domains slightly larger than the trichotomous domain, these axioms are incompatible. Even other strategic properties such as immunity to manipulation via altering of endowments is generally incompatible with individual rationality and Pareto-efficiency [Klaus et al., 2006, Atlamaz and Klaus, 2007]. If we weaken the strategic requirement to say that successful manipulations are computationally intractable to compute, as suggested by Pini et al. [2011], then there are adaptations of the top-trading-cycles algorithm that satisfy this property while maintaining individual rationality and Pareto-efficiency [Fujita et al., 2015, Sikdar et al., 2017, 2018, Phan and Purcell, 2018].

The compatibility of individual rationality, Pareto-efficiency, and strategy-proofness in our model is driven by the restricted domain of preferences. For the two-sided one-to-one matching problem, the analog of this domain—where each agent divides the set partners

⁵In a simple (sequential) priority mechanism, where agents take turns in picking their most preferred bundles without being subjected to individual rationality constraints, an agent’s report can only influence the outcomes for agents who pick after her. It is precisely this feature which makes it straightforward to show that simple priority mechanisms are strategy-proof.

into desirable and undesirable ones and is indifferent among those in the same group—results in these properties being compatible as well [Bogomolnaia and Moulin, 2004]. In the context of balanced exchange in which agents are endowed with and demand multiple objects, Andersson et al. [2018] have independently considered a model similar to ours. By making stronger assumptions on agents’ preferences—that all of the objects that a given agent is endowed with are identical—they are able to design individually rational and strategy-proof mechanisms that are not only Pareto-efficient, but also maximal in the number of objects that are traded. Since there is a trade-off between generality of the model and how demanding an objective can be satisfied, our results are logically independent from theirs.

Biró et al. [2018] consider a model where, as in Andersson et al. [2018], agents are endowed with copies of objects. While they also restrict attention to balanced exchange, they study preferences that are responsive to strict, as opposed to trichotomous, orderings over the objects. They study different variations of the top-trading-cycles algorithm to analyze trade-offs, since in their model Pareto-efficiency and strategy-proofness cannot be reconciled with individual rationality.

Organization of the paper

In Section 2 we introduce the model. In Section 3 we define the IRP algorithm and characterize the outcomes that this algorithm produces. In Section 4 we show that the direct mechanism induced by the IRP algorithm is strategy-proof. In Section 5 we show how to compute IRP matchings in polynomial time. All proofs are in the Appendix.

2 The Model

Let $I = \{1, \dots, n\}$ be a set of n agents and O be a finite set of objects. Each agent $i \in I$ is endowed with a set of objects $\Omega_i \subseteq O$. We assume throughout that $\Omega_i \cap \Omega_j = \emptyset$ whenever $i \neq j$ and that $O = \cup_{i \in I} \Omega_i$.⁶ Each agent $i \in I$ has a weak preference relation \succsim_i over 2^O .

A *matching* is a mapping $\mu : I \rightarrow 2^O$ that satisfies the following two requirements:

1. For any pair of distinct agents $i, j \in I$, $\mu(i) \cap \mu(j) = \emptyset$.
2. For any agent $i \in I$, $|\mu(i)| = |\Omega_i|$.

We require exchange to be *balanced* in the sense that each agent ends up with as many objects as she brings to the market. Let \mathcal{M} be the set of all matchings. Given a profile of preferences

⁶The case where some objects are not in the endowment of any agent, i.e. $\cup_{i \in I} \Omega_i \subsetneq O$, is easy to accommodate by introducing extra (dummy) agents and objects. Details are available upon request.

$\succsim \equiv (\succsim_i)_{i \in I}$ and a matching $\mu \in \mathcal{M}$, μ is *individually rational under* \succsim if each agent finds her assignment to be at least as desirable as her endowment—that is, for each $i \in I$, $\mu(i) \succsim_i \Omega_i$. The matching that assigns each agent her endowment is individually rational. A set of objects $O' \subseteq O$ such that $\Omega_i \succ_i O'$ is *unacceptable* and a set of objects $O'' \subseteq O$ such that $O'' \succeq_i \Omega_i$ is *acceptable* to agent $i \in I$. Next, $\mu' \in \mathcal{M}$ *Pareto-improves* $\mu \in \mathcal{M}$ if, for each $i \in I$, $\mu'(i) \succsim_i \mu(i)$ and, for some $i \in I$, $\mu'(i) \succ_i \mu(i)$. If there is no μ' that Pareto-improves μ , then μ is *Pareto-efficient*.

We now develop assumptions about agents' preferences that play a key role in the remainder of this paper. Throughout, we fix an agent $i \in I$, i 's endowment Ω_i , and a set of *desirable* objects $A_i \subseteq O$ for i . Our first assumption is that, when we restrict attention to acceptable sets of objects, more desirable objects are always better.⁷

Assumption 1 (More is better). For any two sets $B_i, C_i \in 2^O$ such that $B_i \succsim_i \Omega_i$ and $C_i \succsim_i \Omega_i$, $B_i \succ_i C_i$ if and only if $|B_i \cap A_i| > |C_i \cap A_i|$.

Assumption 1 implies that i is indifferent between any two acceptable bundles that contain the same number of desirable objects. Next, we describe two different assumptions about which sets of objects an agent finds acceptable. We show that both assumptions are equivalent when we restrict attention to balanced exchange. Our first assumption describes a scenario in which agent i exhibits an *aversion to unendowed undesirable objects* in the sense that i is not willing to accept a set of objects that contains one or more undesirable objects that she was not endowed with, no matter how many desirable objects the set contains.

Assumption 2 (Aversion to unendowed undesirable objects). For any $B_i \in 2^O$, $B_i \succsim_i \Omega_i$ if and only if $B_i \subseteq A_i \cup \Omega_i$.

In the context of shift exchange, Assumption 2 means that a worker is never willing to create additional scheduling conflicts by accepting a shift for which she already knows that she is not available. By contrast, any shift schedule that leaves i with only desirable shifts and undesirable endowed shifts is deemed acceptable.

Our next assumption describes a scenario in which agent i exhibits an aversion to unfavourable terms of trade in the sense that i requires at least one new desirable object for each of her endowed objects.

Assumption 3 (Aversion to unfavourable terms of trade). For any $B_i \in 2^O$, $B_i \succsim_i \Omega_i$ if and only if $|(B_i \cap A_i) \setminus \Omega_i| \geq |\Omega_i \setminus B_i|$.

⁷In principle, the assumptions that follow allow agents to find it desirable to end up with strictly more objects than they brought to the market. Since we restrict attention to balanced exchange and assume that agents' endowments are known to all, it does not matter for our analysis how an agent ranks bundles with greater cardinality than her endowment.

In the context of shift exchange, Assumption 3 requires that a worker is willing to give up one of her endowed shifts if and only if she gets one new shift for which she has no scheduling conflicts. The assumption is sensible if, for example, i is contractually obliged to work at least $|\Omega_i|$ shifts or needs to work (and be compensated for) at least $|\Omega_i|$ shifts to pay her bills. We now establish that Assumptions 2 and 3 produce exactly the same set of individually rational matchings.

Proposition 1. *Let $i \in I$ be an agent with endowment Ω_i . Assume that both \succsim_i and \succsim'_i satisfy Assumption 1, that \succsim_i satisfies Assumption 2, and that \succsim'_i satisfies Assumption 3.*

Let $B_i \subseteq O$ be such that $|B_i| = |\Omega_i|$. The following are equivalent:

1. B_i is acceptable to i under \succsim_i ,
2. B_i is acceptable to i under \succsim'_i , and
3. $|(B_i \cap A_i) \setminus \Omega_i| = |\Omega_i \setminus B_i|$.

Proposition 1 tells us that if we restrict attention to balanced exchange, then whether we assume that preferences exhibit aversion to unendowed undesirable objects or aversion to unfavourable terms of trade, makes no difference. Hence, we summarize our assumptions on preferences by means of the following definition

Definition 1. Fix an agent $i \in I$, an endowment $\Omega_i \subseteq O$, and a desirable set $A_i \subseteq O$. We say that i 's preferences over sets of objects \succsim_i are *trichotomous* with respect to A_i and Ω_i if \succsim_i satisfies Assumption 1 and either Assumption 2 or Assumption 3.

Before proceeding, we discuss the relationship of our trichotomous preference domain to other approaches in the related literature. For that purpose, note first that we allow for the case where $\Omega_i \cap A_i \notin \{\emptyset, \Omega_i\}$, i.e. the case where some of i 's endowed objects are desirable, while others are undesirable. Hence, we need information about endowments *and* desirable sets of objects in order to describe agents' preferences. To highlight this feature of our model, we have opted for the name trichotomous, as opposed to *dichotomous* as in [Bogomolnaia and Moulin \[2004\]](#).⁸ Next, note that we also allow for the possibility that a non-empty strict subset of j 's endowment Ω_j is desirable for $i \neq j$. This is an important way that ours is different from the the independent contribution of [Andersson et al. \[2018\]](#). There, agents are assumed to either like, or find desirable, none or all of the objects another

⁸[Bogomolnaia and Moulin \[2004\]](#) consider a two-sided one-to-one matching market in which each agent partitions potential match partners from the other side of the market into acceptable (better than being left unmatched) and unacceptable (worse than being left unmatched) ones. Since each agent is endowed with and cannot consume more than one "object", agents' preferences are entirely described by their sets of acceptable partners.

agent brings to the market. While this assumption may be sensible in the context of time banks, which is the leading application in [Andersson et al. \[2018\]](#), it is hard to justify in the context of shift exchange. In the latter application, we can think of objects in i 's desirable set A_i as shifts where i has no scheduling conflicts at all, of objects in $\Omega_i \setminus A_i$ as shifts that i is responsible for but would like to trade away to engage in other activities (e.g. additional paid labor for other firms, further training/education, vacation), and of objects in $O \setminus (A_i \cup \Omega_i)$ as shifts that i is not able to take on due to existing scheduling conflicts. Clearly, it is possible that i is able to take on some shifts initially assigned to j but not others.

Henceforth, we assume that the preferences of all agents are trichotomous. For each agent $i \in I$, the acceptable sets containing $|\Omega_i|$ objects and the ranking of these acceptable sets are then completely identified by her endowment Ω_i and her desirable set A_i . We also assume from now on that endowments are known and fixed at the profile $\Omega = (\Omega_i)_{i \in I}$. In the context of shift exchange, the assumption of known endowments is reasonable since one would expect firms and workers to know who is supposed to work when (if no shift exchanges were to take place) once an initial assignment of shifts has been determined.

Since we assume that preferences are trichotomous and that endowments are fixed and known, the only additional information that we need to know how agents rank individually rational matchings is each agent's desirable set. We assume that this set is private information to be elicited from each agent. In particular, agent i reports an element of $\mathcal{A} \equiv 2^O$ that she claims to be her desirable set. A *mechanism* is a mapping $\varphi : \mathcal{A}^n \rightarrow \mathcal{M}$ that associates a matching to each profile of desirable sets. Given a profile $A \in \mathcal{A}^n$, we denote the set of objects that i receives under mechanism φ by $\varphi_i(A)$. A mechanism is *individually rational* if it selects an individually rational matching for every profile of desirable sets. Recall that our assumptions on preferences imply that μ is individually rational at $A \in \mathcal{A}$ if and only if $\mu(i) \subseteq A_i \cup \Omega_i$. Similarly, a mechanism is *Pareto-efficient* if it selects a Pareto-efficient matching for every profile of desirable sets. Finally, a mechanism is *strategy-proof* if no agent can ever benefit by lying about her desirable set. Our assumptions about preferences imply that an individually rational mechanism φ is strategy-proof if there are no $A \in \mathcal{A}^n$, $i \in I$, and $\hat{A}_i \in \mathcal{A}$ such that $\varphi_i(\hat{A}_i, A_{-i}) \subseteq A_i \cup \Omega_i$ and $|\varphi_i(\hat{A}_i, A_{-i}) \cap A_i| > |\varphi_i(A) \cap A_i|$. Our contribution is to define a mechanism that is individually rational, Pareto-efficient, and strategy-proof on the domain of trichotomous preferences.

3 The Individually Rational Priority Algorithm

In this section, we introduce an algorithm that produces individually rational and Pareto-efficient matchings. Like sequential priority, agents take turns picking their most preferred

sets of objects according to some exogenous priority ranking. The crucial difference with vanilla sequential priority is that our algorithm constrains each agent to choose in a way that is compatible with individual rationality for all agents. Since the set of individually rational matchings varies in response to each agent’s report, an agent can influence the choice set of any other agent—not just those with lower priority. This feature of our algorithm makes it much harder to show that the induced direct mechanism is strategy-proof, which we turn to in the next section, than to show that a simple sequential priority mechanism has this property.

We now describe our algorithm. In the following, we equate an agent’s index $i \in I = \{1, \dots, n\}$ with her priority. Note that lower indices correspond to higher priority. For the remainder of this section, fix a profile of desirable sets $A \in \mathcal{A}$. The *individually rational priority (IRP) algorithm* for A proceeds as follows:

Step 0: Let $\mathcal{M}^0(A)$ be the set of all individually rational matchings at A .

Step $t \in \{1, \dots, n\}$: Let

$$K^t(A) \equiv \max_{\mu \in \mathcal{M}^{t-1}(A)} |\mu(t) \cap A_t|$$

be the *promise to agent t* and

$$\mathcal{M}^t(A) = \{\mu \in \mathcal{M}^{t-1}(A) : |\mu(t) \cap A_t| = K^t(A)\}$$

be the set of all matchings in $\mathcal{M}^{t-1}(A)$ that comply with the promise to t .

The above definitions imply that for any pair t, t' such that $t' \leq t$ and any $\mu \in \mathcal{M}^t(A)$, $|\mu(t') \cap A_{t'}| = K^{t'}(A)$. In particular, all matchings in $\mathcal{M}^n(A)$ are welfare equivalent for all agents. We call any $\mu \in \mathcal{M}^n(A)$ an *IRP outcome* for A .

We now establish that all IRP outcomes are individually rational and Pareto-efficient. First, for any $t \in \{1, \dots, n\}$, we have $\mathcal{M}^{t-1}(A) \subseteq \mathcal{M}^t(A)$. Hence, $\mathcal{M}^n(A) \subseteq \mathcal{M}^0(A)$, which implies that IRP outcomes are individually rational. Second, let $\mu \in \mathcal{M}^n(A)$. The construction of the sequence $\{\mathcal{M}^t(A)\}_{t=1}^n$ implies that there is no matching in $\mathcal{M}^0(A)$ that Pareto-dominates μ . For any matching $\nu \in \mathcal{M} \setminus \mathcal{M}^0(A)$, there is at least one agent $j \in I$ for whom individual rationality is violated, i.e. $\mu(j) \succ_j \Omega_j \succ_j \nu(j)$. Hence, there is no matching in \mathcal{M} that Pareto-dominates μ . The following result summarizes the findings of this paragraph.

Theorem 1. *Any $\mu \in \mathcal{M}^n(A)$ is individually rational and Pareto-efficient.*

In the remainder of this section, we develop a characterization of matchings in the sequence $\{\mathcal{M}^t(A)\}_{t=1}^n$ that plays a crucial role for our subsequent analysis. For this purpose, we introduce some additional definitions.

Given a matching $\mu \in \mathcal{M}^0(A)$, a *cycle of μ* is a sequence $C = (i^1, o^1, \dots, i^M, o^M)$ of M distinct agents and M distinct objects such that, for each $m \in \{1, \dots, M\}$, $o^{m-1} \in \mu(i^m)$ and $o^m \notin \mu(i^m)$, where $o^0 \equiv o^M$. For the following discussion, fix an arbitrary cycle $C = (i^1, o^1, \dots, i^M, o^M)$ of μ . For any $m \in \{1, \dots, M\}$, we say that i^m *points to* o^m and that o^m *points to* i^{m+1} , where $i^{M+1} \equiv i^1$. Thus, C is a sequence of agent-object pairs such that each agent i^m points to an object o^m that she does not get at μ , i.e. an object $o^m \notin \mu(i^m)$, and each object o^m points to the agent who gets it at μ , i.e. the agent i^{m+1} for whom $o^m \in \mu(i^{m+1})$. Let $I(C) \equiv \{i^1, \dots, i^M\}$ be the set of agents involved in C and let $O(C) \equiv \{o^1, \dots, o^M\}$ be the set of objects involved in C . Let $\mu + C$ denote the matching that results from μ by executing the trades designated by C , i.e., for each $k \in I$,

$$(\mu + C)(k) \equiv \begin{cases} \mu(k) & \text{if } k \notin I(C) \\ (\mu(k) \setminus \{o^{m-1} : i^m = k\}) \cup \{o^m : i^m = k\} & \text{if } k \in I(C). \end{cases}$$

Motivated by the definition of $\mu + C$, we say that, for any $m \in \{1, \dots, M\}$, i^m *trades* o^{m-1} for o^m in C . We say that C

- is *individually rational* if, for each $m \in \{1, \dots, M\}$, $o^m \in \Omega_{i^m} \cup A_{i^m}$,
- *increases (decreases) i 's welfare* if $|(\mu + C)(i) \cap A_i| > (<) |\mu(i) \cap A_i|$, and
- *affects i* if it either increases or decreases i 's welfare.

Finally, we say that C is a *constrained improvement cycle (CIC) of μ for t at A* if C

1. individually rational,
2. increases t 's welfare, and
3. does not affect any agent $t' < t$.

With these definitions, we now state the second result of this section.

Theorem 2. *For any $t \in \{1, \dots, n\}$, we have that*

- (i) $\mu \in \mathcal{M}^t(A)$ only if there does not exist a CIC of μ for some $t' \leq t$ at A , and
- (ii) if $\mu \in \mathcal{M}^{t-1}(A)$ and there does not exist a CIC of μ for t at A , then $\mu \in \mathcal{M}^t(A)$.

To gain some intuition for Theorem 2, consider the t^{th} step of the IRP algorithm. Here, the number of desirable objects awarded to agent t is determined subject to the constraints imposed by the promises to agents $t' < t$. These constraints are expressed in the restriction to matchings in $\mathcal{M}^{t-1}(A)$. Now fix an arbitrary matching $\mu \in \mathcal{M}^t(A)$. If C is a CIC of μ for $t' \leq t$ at A , then $\mu + C \in \mathcal{M}^{t'-1}(A)$ given that $\mu \in \mathcal{M}^t(A) \subseteq \mathcal{M}^{t'-1}(A)$. Since C increases the welfare of t' , we find that $\mu \notin \mathcal{M}^{t'}(A)$ and hence, given that $t' \leq t$, $\mu \notin \mathcal{M}^t(A)$. For the proof of part (ii), pick an arbitrary $\nu \in \mathcal{M}^{t-1}(A)$ that leaves t strictly better off than μ . Then there exists an object o that is desirable for t and that t gets in ν but not in μ . In our proof, we use o as well as the other assignments in μ and ν to construct a CIC of μ for j at A .

4 IRP Mechanisms and Strategy-proofness

In this section, we show that a direct mechanism induced by the IRP algorithm for any given priority order is strategy-proof. Mechanism φ is an *individually rational priority (IRP) mechanism* if $\varphi(A) \in \mathcal{M}^n(A)$ for all $A \in \mathcal{A}$. Recall that, since we have fixed the order over agents, any IRP mechanism is guaranteed to induce the same welfare for all agents. Recall also that $\{\mathcal{M}^t(A)\}_{t=1}^n$ summarizes the progression of the IRP algorithm for input A .

For the remainder of this section, fix a true profile of desirable sets $A \in \mathcal{A}$, an agent $i \in I$, and a possible manipulation for i , $\hat{A}_i \in \mathcal{A}_i$. Denote the profile (\hat{A}_i, A_{-i}) by \hat{A} . We establish that an IRP mechanism is strategy-proof by showing that, when the true profile of desirable sets is A , i is no better off at \hat{A} than he is at A .

We start by showing that i does not benefit by falsely claiming that an object is desirable when it is not. The first of two lemmas considers the case where she falsely claims as desirable an object o that she is not endowed with. It says that the the number of trades that she receives can only change if she receives o at every matching produced by the IRP algorithm at \hat{A} .

Lemma 1. *If, for some $o \notin \Omega_i \cup A_i$, $\hat{A}_i = A_i \cup \{o\}$ and $K^i(\hat{A}) \neq K^i(A)$, then, for each $\mu \in \mathcal{M}^n(\hat{A})$, $o \in \mu(i)$.*

Since preferences are trichotomous, Lemma 1 implies that falsely claiming a single unendowed undesirable object to be desirable is not beneficial. The next lemma shows that an agent cannot benefit from falsely claiming that one of her endowed objects is desirable.

Lemma 2. *If, for some $o \in \Omega_i \setminus A_i$, $\hat{A}_i = A_i \cup \{o\}$, then for each $\mu \in \mathcal{M}^i(\hat{A})$, $K^i(A) \geq |\mu(i) \cap A_i|$.*

Lemmas 1 and 2 imply that expanding the set of desirable objects can never be profitable for an agent. Next, we consider the case where i falsely declares a desirable object to be undesirable.

Lemma 3. *If, for some $o \in A_i$, $\hat{A}_i = A_i \setminus \{o\}$, then $K^i(A) \geq K^i(\hat{A})$.*

Since Lemma 3 is the main step in proving strategy-proofness of IRP mechanisms, we provide a sketch of its proof. Note first that when i shrinks her desirable set from A_i to $\hat{A}_i = A_i \setminus \{o\}$, she places an additional constraint on the choices of all agents in the IRP algorithm since she can no longer receive o . Put differently, the set of individually rational matchings shrinks from $\mathcal{M}^0(A)$ to $\mathcal{M}^0(\hat{A}) \subseteq \mathcal{M}^0(A)$. It is then immediate that a contraction can never be profitable for the highest priority agent since such a manipulation would only shrink her own choice set. Things are potentially different for $i > 1$: she might hope that shrinking the choice sets of agents $j < i$ forces these agents to leave her with a choice set $\mathcal{M}^{i-1}(\hat{A})$ that contains some matching μ that she prefers to all matchings in $\mathcal{M}^{i-1}(A)$. So assume that there exists such a matching $\mu \in \mathcal{M}^{i-1}(\hat{A})$. Then $|\mu(i) \cap A_i| > K^i(A)$. Since $\mu \notin \mathcal{M}^{i-1}(A)$ but $\mu \in \mathcal{M}^0(A)$, there exists a smallest integer $j < i$ such that $\mu \notin \mathcal{M}^j(A)$. By Theorem 2, there exists a CIC C of μ for j at A . Given that the only difference between A and \hat{A} is that i ranks o as undesirable in \hat{A} , i points to o in C . Since $o \in A_i \setminus \Omega_i$, C cannot decrease the welfare of i . If $\mu + C \in \mathcal{M}^{i-1}(A)$, we obtain a contradiction to our assumption that $|\mu(i) \cap A_i| > K^i(A)$ since there is a matching that i could have picked in the IRP algorithm at A that is at least as good as μ . If $\mu + C \notin \mathcal{M}^{i-1}(A)$, Theorem 2 again implies that there has to be a CIC C' of $\mu + C$ at A for some $j' < i$. The main part of our proof of Lemma 3 shows that we can construct a third CIC C^* from C and C' such that $\mu + C^*$ (A) makes every agent $k \leq \min\{j, j'\}$ at least as well off as $(\mu + C) + C'$, and (B) makes i weakly better off than μ . Iterating this argument, we eventually obtain a contradiction to our assumption that $|\mu(i) \cap A_i| > K^i(A)$ since there exists at least one matching in $\mathcal{M}^{i-1}(A)$ that makes i weakly better off than μ .

Finally, we appeal to Lemmas 1, 2, and 3 to prove that any IRP mechanism is strategy-proof.

Theorem 3. *Any IRP mechanism is strategy-proof.*

We point out that our restriction to trichotomous preferences is important for strategy-proofness of IRP mechanisms. For example, even for the case of singleton endowments, a sequential priority over the set of individually rational matchings is not generally strategy-proof if agents' preferences have more than three indifference classes.

5 Computing IRP Matchings

In this section, we define a polynomial time algorithm (Algorithm 1) to compute matchings in $\mathcal{M}^n(A)$ based on network flows. The procedure described in Section 3 relies on an enumeration of all of the individually rational matchings ($\mathcal{M}^0(A)$), the subset of these matchings in which 1 receives the most desirable objects ($\mathcal{M}^1(A)$), and so on. The problem is that sizes of these sets are an exponential function of the number of agents and objects. The algorithm that we now develop uses the fact that each of these sets is the set of solutions to a particular network flow problem.

For the following discussion, fix an endowment Ω and a profile of desirable sets A . We work with a directed graph (V, E) . The set of vertices V contains three vertices corresponding to each $i \in I$, call them i, i^A , and i^U . It also contains one vertex corresponding to each $o \in O$. Finally, it contains a source S and a sink T . That is,

$$V \equiv \left[\bigcup_{i \in I} \{i, i^A, i^U\} \right] \cup O \cup \{S, T\}$$

The set of edges E contains an edge from S to each $i \in I$, from each $i \in I$ to i^A and i^U , from i^A to each object in A_i , from i^U to each object in $\Omega_i \setminus A_i$, and from each $o \in O$ to T . That is,

$$\begin{aligned} E &\equiv \{(S, i) : i \in I\} \\ &\cup \{(i, i^A), (i, i^U) : i \in I\} \\ &\cup \{(i^A, o) : o \in A_i\} \\ &\cup \{(i^U, o) : o \in \Omega_i \setminus A_i\} \\ &\cup \{(o, T) : o \in O\}. \end{aligned}$$

Roughly speaking, the idea here is that copy i^A of agent i collects all desirable objects, while copy i^U collects all undesirable endowments. Throughout our algorithm, we only vary the capacities of each edge. The capacity of edge $e \in E$ is $q(e)$ and we initialize q as follows:

$$\begin{aligned} q(S, i) &= |\Omega_i| & i \in I \\ q(i, i^A) &= |\Omega_i| & i \in I \\ q(i, i^U) &= |\Omega_i \setminus A_i| & i \in I \\ q(i^A, o) &= 1 & i \in I, o \in A_i \\ q(i^U, o) &= 1 & i \in I, o \in \Omega_i \\ q(o, T) &= 1 & o \in O. \end{aligned}$$

We say that $f : E \rightarrow \mathbb{N}$ is an (integer) flow, if $f(e) \leq q(e)$, for all $e \in E$, and

$\sum_{e \in \delta^-(v)} f(e) = \sum_{e \in \delta^+(v)} f(e)$, for all $v \in V \setminus \{S, T\}$.⁹ The value of a flow f through i is $f(S, i)$ and the value of a flow f is $v(f) = \sum_{i \in I} f(S, i)$. Since $q(S, i) = |\Omega_i|$, for all $i \in I$, the maximum flow through i cannot be larger than $|\Omega_i|$. Since for all $o \in O$, $i \in I$, and $D \in \{A, U\}$, $q(i^D, o) \in \{0, 1\}$, a flow cannot use two edges involving the same object but different agents. If no flow g has higher value than f , then f is a *maximum flow*. Let $\text{MAXFLOW}(q)$ be the value of a maximum flow from S to T when the capacities are q . By the above observation that the flow through i cannot exceed $|\Omega_i|$, for every q that we consider, $\text{MAXFLOW}(q) \leq \sum_{i \in I} |\Omega_i|$ and, for the initial value of q , $\text{MAXFLOW}(q) = \sum_{i \in I} |\Omega_i|$ since we can assign each agent her endowment at that capacity vector. Given $\mu \in \mathcal{M}$, we say that μ *corresponds to* f if, for each $i \in I$ and each $o \in \mu(i) \cap A_i$, $f(i^A, o) = 1$ and, for each $i \in I$ and each $o \in \mu(i) \cap \Omega_i \setminus A_i$, $f(i^u, o) = 1$.

Algorithm 1 iteratively minimizes each agent i 's *undesirable object capacity*, $q(i, i^U)$, by solving a series of network flow problems.

Algorithm 1 Procedure to compute IRP

```

1: procedure IRP( $V, E, q$ )
2:   for  $t = 1$  to  $t = N$  do                                ▷ Loop invariant:  $\text{MAXFLOW}(q) = \sum_i |\Omega_i|$ 
3:     if  $q(t, t^U) > 0$  then                                ▷  $t$  has undesirable object capacity
4:        $\hat{q} = q$ 
5:        $\hat{q}(t, t^U) = \hat{q}(t, t^U) - 1$                         ▷ Decrement  $t$ 's undesirable capacity by 1
6:       while  $\text{MAXFLOW}(\hat{q}) = \sum_i |\Omega_i|$  do          ▷ As long as the loop invariant holds
7:          $q = \hat{q}$ 
8:          $\hat{q}(t, t^U) = \hat{q}(t, t^U) - 1$                     ▷ Decrement  $t$ 's undesirable capacity by 1
9:   return  $q$ 

```

The following result summarizes the key properties of Algorithm 1.

Theorem 4. *Let $A \in \mathcal{A}$ be a profile of desirable sets and let Ω be an endowment. Suppose that V, E , and q are defined based on A and Ω as described above.*

1. *A flow f is a maximum flow in $(V, E, \text{IRP}(V, E, q))$ if and only if it corresponds to an IRP outcome.*
2. *Fixing Ω , the complexity of Algorithm 1 is $O(mn^3)$, where $m = |O|$.*

6 Conclusion

We have considered a model of exchange of indivisible goods without monetary transfers and multi-unit demand and supply. The following corollary summarizes our main results.

⁹By $\delta^+(v)$ we denote the edges that originate at node v and by $\delta^-(v)$ the edges that point to v .

Corollary 1. *Fix an endowment Ω and assume that all agents have trichotomous preferences.*

1. *IRP mechanisms are individually rational, Pareto-efficient, and strategy-proof.*
2. *It is computationally efficient to find IRP outcomes.*

In light of Corollary 1, IRP mechanisms are promising solutions to balanced exchange problems in which agents have trichotomous preferences and are endowed with more than one indivisible object.

References

- M. Akbarpour, S. Li, and S. O. Gharan. Thickness and information in dynamic matching markets. Working paper, 2018.
- J. Alcalde-Unzu and E. Molis. Exchange of indivisible goods and indifferences: The top trading absorbing sets mechanisms. *Games and Economic Behavior*, 73(1):1–16, September 2011.
- T. Andersson, Ágnes Cseh, L. Ehlers, and A. Erlanson. Organizing time banks: Lessons from matching markets. Working paper, 2018.
- M. Atlamaz and B. Klaus. Manipulation via Endowments in Exchange Markets with Indivisible Goods. *Social Choice and Welfare*, 28(1):1–18, January 2007.
- H. Aziz and B. De Keijzer. Housing markets with indifferences: A tale of two mechanisms. In *AAAI*, volume 12, pages 1–1, 2012.
- P. Biró, F. Klijn, and S. Pàpai. Balanced exchange in a multi-object shapley-scarf market. Working paper, 2018.
- A. Bogomolnaia and H. Moulin. Random matching under dichotomous preferences. *Econometrica*, 72(1):257–279, 01 2004.
- E. Fujita, J. Lesca, A. Sonoda, T. Todo, and M. Yokoo. A complexity approach for core-selecting exchange with multiple indivisible goods under lexicographic preferences. In *Proceedings of the Twenty-Ninth AAAI Conference on Artificial Intelligence, AAAI’15*, pages 907–913, 2015. ISBN 0-262-51129-0.
- P. Jaramillo and V. Manjunath. The difference indifference makes in strategy-proof allocation of objects. *Journal of Economic Theory*, 147(5):1913–1946, 2012.

- B. Klaus, D. Dimitrov, and C.-J. Haake. Bundling in exchange markets with indivisible goods. *Economics Letters*, 93(1):106–110, 2006.
- V. Manjunath and A. Westkamp. Strategy-proof exchange beyond trichotomous preferences. Working paper, 2018.
- W. Phan and C. Purcell. On the complexity of manipulation in the exchange of indivisible objects. Working paper, 2018.
- M. S. Pini, F. Rossi, K. B. Venable, and T. Walsh. Manipulation complexity and gender neutrality in stable marriage procedures. *Autonomous Agents and Multi-Agent Systems*, 22(1):183–199, Jan 2011.
- D. Saban and J. Sethuraman. House allocation with indifferences: A generalization and a unified view. In *Proceedings of the Fourteenth ACM Conference on Electronic Commerce*, EC '13, pages 803–820, 2013. ISBN 978-1-4503-1962-1.
- L. Shapley and H. Scarf. On cores and indivisibility. *Journal of Mathematical Economics*, 1: 23–37, 1974.
- S. Sikdar, S. Adali, and L. Xia. Mechanism design for multi-type housing markets. In *Proceedings of the Thirty-First AAAI Conference on Artificial Intelligence*, AAAI'17, pages 684–690, 2017.
- S. Sikdar, S. Adali, and L. Xia. Top-trading-cycles mechanisms with acceptable bundles. In *Proceedings of the Thirty-Second AAAI Conference on Artificial Intelligence*, AAAI'18, 2018.
- T. Sönmez. Strategy-proofness and essentially single-valued cores. *Econometrica*, 67(3): 677–690, May 1999.

Appendices

A Proof of Proposition 1

Since $|B_i| = |\Omega_i|$,

$$|B_i \setminus \Omega_i| = |\Omega_i \setminus B_i|. \tag{1}$$

Furthermore, since $[(B_i \cap A_i) \setminus \Omega_i] \cup (B_i \cap \Omega_i) \subseteq B_i$, we have

$$|(B_i \cap A_i) \setminus \Omega_i| + |B_i \cap \Omega_i| \leq |B_i|. \quad (2)$$

Since $\Omega_i = (\Omega_i \setminus B_i) \cup (B_i \cap \Omega_i)$, we have $|\Omega_i| = |\Omega_i \setminus B_i| + |B_i \cap \Omega_i|$. Thus, using $|B_i| = |\Omega_i|$, (2) implies

$$|(B_i \cap A_i) \setminus \Omega_i| \leq |\Omega_i \setminus B_i|. \quad (3)$$

By Assumption 2, B_i is acceptable to i under \succsim_i if and only if $B_i \subseteq A_i \cup \Omega_i$, or equivalently $|(B_i \cap A_i) \setminus \Omega_i| = |B_i \setminus \Omega_i|$. By (1) this is equivalent to $|(B_i \cap A_i) \setminus \Omega_i| = |\Omega_i \setminus B_i|$

By Assumption 3, B_i is acceptable to i under \succsim'_i if and only if $|(B_i \cap A_i) \setminus \Omega_i| \geq |\Omega_i \setminus B_i|$, or equivalently by (3), $|(B_i \cap A_i) \setminus \Omega_i| = |\Omega_i \setminus B_i|$.

B Proof of Theorem 2

We first introduce a notion of generalized CICs where we allow agents, but not objects, to reoccur. A *generalized cycle* of μ is a sequence $C \equiv (i^1, o^1, \dots, i^M, o^M)$ of M (not necessarily distinct) agents and M distinct objects such that, for each $m \in \{1, \dots, M\}$, $o^{m-1} \in \mu(i^m)$ and $o^m \notin \mu(i^m)$ (where $o^0 \equiv o^M$). For the following discussion, fix an arbitrary generalized cycle $C = (i^1, o^1, \dots, i^M, o^M)$ of μ . As for the case of cycles, we denote the matching that results by implementing the exchanges that C specifies by $\mu + C$, i.e., for each $k \in I$, we set

$$(\mu + C)(k) \equiv \begin{cases} \mu(k) & \text{if } k \notin I(C) \\ (\mu(k) \setminus \{o^{m-1} : i^m = k\}) \cup \{o^m : i^m = k\} & \text{if } k \in I(C) \end{cases}$$

We say that C is an *individually rational generalized cycle*, if, for all m , $o^m \in \Omega_{i^m} \cup A_{i^m}$. Furthermore, we refer to i^1 as the *head* of C . A *generalized constrained improvement cycle (GCIC) of μ for j at A* is an individually rational generalized cycle $C = (i^1, o^1, \dots, i^M, o^M)$ of μ at A that satisfies the following additional conditions:

1. j is the head of C : $i^1 = j$,
2. C improves its head's welfare: $o^M \in \Omega_j \setminus A_j$, $o^1 \in A_j$, and, for any $m \neq 1$ such that $i^m = j$ and $o^m \in \Omega_j \setminus A_j$, we have that $o^{m-1} \in \Omega_j \setminus A_j$
3. C does not affect agents with higher priority than its head: for any m such that $i^m < j$, either $\{o^{m-1}, o^m\} \subseteq A_{i^m}$ or $\{o^{m-1}, o^m\} \subseteq \Omega_{i^m} \setminus A_{i^m}$.

We now start our proof by verifying that if C is an individually rational generalized cycle, then $\mu + C$ is individually rational as well.

Lemma 4. *Let $A \in \mathcal{A}$ and $\mu \in \mathcal{M}^0(A)$. If C is an individually rational generalized cycle of μ at A , then $\mu + C \in \mathcal{M}^0(A)$.*

Proof. Let $C = (i^1, o^1, \dots, i^M, o^M)$ and $\nu \equiv \mu + C$. First, for each $o \in O$, since the objects in C are distinct o cannot be in both $\nu(k)$ and $\nu(l)$ for distinct agents k and l . Next, for each $k \in I$, $\nu(k) \subseteq \Omega_k \cup A_k$ since $\mu(k) \subseteq \Omega_k \cup A_k$ and $\nu(k) \setminus \mu(k) \subseteq \Omega_k \cup A_k$ given that C is individually rational. Finally, we verify that for each $k \in I$, $|\nu(k)| = |\Omega_k|$. Since the objects in C are distinct and since $o^{m-1} \in \mu(i^m)$ as well as $o^m \notin \mu(i^m)$ for all m , we have

$$\begin{aligned} |\nu(k)| &= |\mu(k) \setminus \{o^{m-1} : i^m = k\}| + |\{o^m : i^m = k\}| \\ &= |\mu(k)| - |\{o^{m-1} : i^m = k\}| + |\{o^m : i^m = k\}| \\ &= |\Omega_k| \end{aligned}$$

□

For the remainder of the proof, we fix $t \in \{1, \dots, n\}$. We first argue that the absence of GCICs is necessary for a matching to be compatible with the guarantee for t at A .

Lemma 5. *If $\mu \in \mathcal{M}^t(A)$, then there is no GCIC of μ for any $t' \leq t$ at A .*

Proof. Suppose to the contrary that $C \equiv (i^1, o^1, \dots, i^M, o^M)$ is a GCIC of μ for some $t' \leq t$ at A . Let $\nu \equiv \mu + C$. Since C does not affect any agent $t'' < t'$, $|\nu(t'') \cap A_{t''}| = |\mu(t'') \cap A_{t''}| = K^{t''}(A)$ for all $t'' < t'$. By Lemma 4, ν is individually rational. Thus, combining the last two observations with the fact that $\mu \in \mathcal{M}^t(A) \subseteq \mathcal{M}^{t'-1}(A)$, $\nu \in \mathcal{M}^{t'-1}(A)$. However, since C increases t' 's welfare, we have $|\nu(t') \cap A_{t'}| > |\mu(t') \cap A_{t'}| = K^{t'}(A)$ and thus $\mu \notin \mathcal{M}^{t'-1}(A)$. Hence, we obtain a contradiction to $\mu \in \mathcal{M}^t(A)$. □

Since any CIC is, trivially, also a GCIC, Lemma 5 already shows the necessity part, part (i), of Theorem 2.

Showing that the absence of CICs is sufficient, i.e. part (ii) of Theorem 2, requires more work. We begin with an auxiliary lemma showing that while an agent $j < t$ may appear more than once in a GCIC of μ , it has to be the case that j either only trades desirable for desirable objects, or only trades undesirable endowed objects for undesirable endowed objects.

Lemma 6. *Let $j \in I$ be such that $j < t$, $\mu \in \mathcal{M}^{t-1}(A)$, and $C = (i^1, o^1, \dots, i^M, o^M)$ be a GCIC of μ for t at A . If m and m' are such that $i^m = i^{m'} = j$, then either*

$$\begin{aligned} \{o^{m-1}, o^m, o^{m'-1}, o^{m'}\} &\subseteq \Omega_j \setminus A_j \\ &\text{or} \\ \{o^{m-1}, o^m, o^{m'-1}, o^{m'}\} &\subseteq A_j. \end{aligned}$$

Proof. Fix m and m' such that $i^m = i^{m'} = j$ and assume, without loss of generality, that $m < m' \leq M$. Since C does not affect any agent with higher priority than t , it suffices to show that either $\{o^m, o^{m'}\} \subseteq \Omega_j \setminus A_j$ or $\{o^m, o^{m'}\} \subseteq A_j$. Suppose, for the sake of contradiction, that neither is true. There are two cases. If $o^m \in \Omega_j \setminus A_j$ and $o^{m'} \in A_j$, consider

$$C' \equiv (i^1, o^1, \dots, i^{m-1}, o^{m-1}, i^{m'}, o^{m'}, \dots, i^M, o^M).$$

Since C is a generalized individually rational cycle of μ , C' is a generalized individually rational cycle of μ as well. Since C is a GCIC for $t > j$ and $o^m \in \Omega_j \setminus A_j$, we have that $o^{m-1} \in \Omega_j \setminus A_j$ as well. Furthermore, since C is a GCIC for $t > j$, we also obtain that, for any $\tilde{m} \in \{1, \dots, m-1, m'+1, \dots, M\}$ such that $i^{\tilde{m}} \leq t-1$, $\{o^{\tilde{m}-1}, o^{\tilde{m}}\} \subseteq A_{i^{\tilde{m}}}$ or $\{o^{\tilde{m}-1}, o^{\tilde{m}}\} \subseteq \Omega_{i^{\tilde{m}}} \setminus A_{i^{\tilde{m}}}$ (where $o^0 = o^M$). Hence, subject to relabeling the agents, C' is a GCIC of μ for j at A , which we have already argued to be impossible. On the other hand, if $o^m \in A_j$ and $o^{m'} \in \Omega_j \setminus A_j$, the following sequence yields a similar contradiction:

$$(i^m, o^m, \dots, i^{m'-1}, o^{m'-1}).$$

Thus, either $\{o^m, o^{m'}\} \subseteq A_j$ or $\{o^m, o^{m'}\} \subseteq \Omega_j \setminus A_j$. \square

Building on Lemma 6, our next auxiliary lemma states that it is sufficient to consider CICs since any recurrence of an agent can be eliminated to yield a shorter cycle.

Lemma 7. *Let $\mu \in \mathcal{M}^{t-1}(A)$. If there exists a GCIC of μ for t at A , then there exists an CIC of μ for t at A .*

Proof. Let $C \equiv (i^1, o^1, \dots, i^M, o^M)$ be a GCIC of μ for t at A . We show how to construct a CIC C' of μ for t on basis of C .

We first eliminate duplicates of any $j \in I(C)$ such that $t < j$. Suppose there are two indices m and m' such that $m < m'$ and $i^m = i^{m'} = j$. Then, since $\{o^{m-1}, o^m, o^{m'-1}, o^{m'}\} \subseteq \Omega_j \cup A_j$, the sequence $(i^1, o^1, \dots, i^{m-1}, o^{m-1}, i^{m'}, o^{m'}, \dots, i^M, o^M)$ is also a GCIC of μ for t at A that has one fewer instances of j than C .

Next, we eliminate duplicates of t . Suppose there is m such that $m > 1$ and $i^m = t$. If $o^m \in \Omega_t \setminus A_t$, then since C improves its head's welfare, we have $o^{m-1} \in \Omega_t \setminus A_t$, so that $(i^1, o^1, \dots, i^{m-1}, o^{m-1})$ is a GCIC of μ for t at A with one fewer instances of t than C . If $o^m \in A_t$, then $(i^m, o^m, \dots, i^M, o^M)$ is a GCIC of μ for t at A with one fewer instances of t .

Finally, we eliminate duplicates of $j \in I(C)$ such that $j < t$. Suppose there are two indices m and m' such that $m < m'$ and $i^m = i^{m'} = j$. By Lemma 6, we have that either $\{o^{m-1}, o^{m'}\} \subseteq \Omega_j \setminus A_j$ or $\{o^{m-1}, o^{m'}\} \subseteq A_j$. Hence, the sequence $(i^1, o^1, \dots, i^{m-1}, o^{m-1}, i^{m'}, o^{m'}, \dots, i^M, o^M)$ is a GCIC of μ for t at A that has one fewer instances of j than C . \square

In the remainder of this proof, we establish that if $\mu \in \mathcal{M}^{t-1}(A)$ is such that $|\mu(t) \cap A_t| < K^t(A)$, then there is a GCIC of μ for t at A . By Lemma 7, we thereby establish the sufficiency part of Theorem 2.

Fix $\mu \in \mathcal{M}^{t-1}(A)$ such that $|\mu(t) \cap A_t| < K^t(A)$. We construct a GCIC of μ for t at A . We start with $i^1 = t$. Let $\nu \in \mathcal{M}^t(A)$. Since $|\mu(t) \cap A_t| < K^t(A) = |\nu(t) \cap A_t|$, there exists $o^1 \in (\nu(t) \setminus \mu(t)) \cap A_t$. Suppose now that we have grown a sequence $(i^1, o^1, \dots, i^M, o^M)$ of M agents and distinct objects in a way that satisfies the following four conditions:

$$\text{for each } m \geq 1, o^m \in \nu(i^m) \setminus \mu(i^m), \text{ and, for each } m \geq 2, o^{m-1} \in \mu(i^m) \setminus \nu(i^m), \quad (\text{C1})$$

$$\text{for each } m \geq 1, o^m \in \Omega_{i^m} \cup A_{i^m}, \quad (\text{C2})$$

$$\begin{aligned} &\text{for each pair } m, m' \text{ such that } i^m = i^{m'} < t, \text{ either} \\ &\{o^{m-1}, o^m, o^{m'-1}, o^{m'}\} \subseteq A_{i^m} \text{ or } \{o^{m-1}, o^m, o^{m'-1}, o^{m'}\} \subseteq \Omega_{i^m} \setminus A_{i^m}, \text{ and} \end{aligned} \quad (\text{C3})$$

$$\text{for each } m \geq 2 \text{ such that } i^m = t, \{o^{m-1}, o^m\} \subseteq A_t. \quad (\text{C4})$$

(C1) to (C4) imply that if $o^M \in \mu(i^1) \cap (\Omega_{i^1} \setminus A_{i^1})$, then $(i^1, o^1, \dots, i^M, o^M)$ is a GCIC of μ for $i^1 = t$ at A . We now show that if $o^M \notin \mu(i^1) \cap (\Omega_{i^1} \setminus A_{i^1})$, then we can find an agent-object pair (i^{M+1}, o^{M+1}) so that $(i^1, o^1, \dots, i^{M+1}, o^{M+1})$ satisfies (C1) to (C4).

Since both ν and μ are matchings, $\sum_{i \in I} |\nu(i)| = \sum_{i \in I} |\mu(i)| = \sum_{i \in I} |\Omega_i|$. Hence, $o^M \notin \mu(i^M)$ implies that there exists an agent $i^{M+1} \in I \setminus \{i^M\}$ such that $o^M \in \mu(i^{M+1})$. Since $o^M \in \nu(i^M)$, we have $o^M \in \mu(i^{M+1}) \setminus \nu(i^{M+1})$. We distinguish three cases according to the identity of i^{M+1} . In each of these cases, we identify an object o^{M+1} such that the sequence $(i^1, o^1, \dots, i^{M+1}, o^{M+1})$ satisfies (C1) to (C4).

Case 1: $i^{M+1} \notin \{i^1, \dots, i^M\}$ There are three subcases to consider:

Case 1.1: $i^{M+1} < t$ and $o^M \in A_{i^{M+1}}$ Since $\nu, \mu \in \mathcal{M}^{t-1}(A)$ and $i^{M+1} < t$, it follows that $|\nu(i^{M+1}) \cap A_{i^{M+1}}| = |\mu(i^{M+1}) \cap A_{i^{M+1}}|$. Since $o^M \in (\mu(i^{M+1}) \setminus \nu(i^{M+1})) \cap A_{i^{M+1}}$ there is $o^{M+1} \in (\nu(i^{M+1}) \setminus \mu(i^{M+1})) \cap A_{i^{M+1}}$. Furthermore, since $i^{M+1} \notin \{i^1, \dots, i^M\}$ and, for each m , $o^m \in \nu(i^m)$, we have that $o^{M+1} \notin \{o^1, \dots, o^M\}$.

Case 1.2: $i^{M+1} < t$ and $o^M \in \Omega_{i^{M+1}} \setminus A_{i^{M+1}}$ Since $\nu, \mu \in \mathcal{M}^{t-1}(A)$ and $i^{M+1} < t$, we have $|\nu(i^{M+1}) \cap (\Omega_{i^{M+1}} \setminus A_{i^{M+1}})| = |\mu(i^{M+1}) \cap (\Omega_{i^{M+1}} \setminus A_{i^{M+1}})|$. Since $o^M \in (\mu(i^{M+1}) \setminus \nu(i^{M+1})) \cap (\Omega_{i^{M+1}} \setminus A_{i^{M+1}})$ there is $o^{M+1} \in (\nu(i^{M+1}) \setminus \mu(i^{M+1})) \cap (\Omega_{i^{M+1}} \setminus A_{i^{M+1}})$. Furthermore, since $i^{M+1} \notin \{i^1, \dots, i^M\}$ and, for each m , $o^m \in \nu(i^m)$, we have that $o^{M+1} \notin \{o^1, \dots, o^M\}$.

Case 1.3: $i^{M+1} > t$ Since ν and μ are individually rational matchings, $\nu(i^{M+1}) \subseteq \Omega_{i^{M+1}} \cup A_{i^{M+1}}$, $\mu(i^{M+1}) \subseteq \Omega_{i^{M+1}} \cup A_{i^{M+1}}$, and $|\nu(i^{M+1})| = |\mu(i^{M+1})| = |\Omega_{i^{M+1}}|$.

Given these facts and that $o^M \in \mu(i^{M+1}) \setminus \nu(i^{M+1})$, there is $o^{M+1} \in \nu(i^{M+1}) \setminus \mu(i^{M+1})$. Furthermore, since $i^{M+1} \notin \{i^1, \dots, i^M\}$ and for each m , $o^m \in \nu(i^m)$, we have that $o^{M+1} \notin \{o^1, \dots, o^M\}$.

Case 2: $i^{M+1} \in \{i^2, \dots, i^M\} \setminus \{t\}$ Let $m \leq M$ be such that $i^m = i^{M+1}$. We distinguish three subcases parallel to those of Case 1.

Case 2.1: $i^{M+1} < t$ and $o^M \in A_{i^{M+1}}$ We first show that $o^m \in A_{i^{M+1}}$. Otherwise, $o^m \in \Omega_{i^{M+1}} \setminus A_{i^{M+1}}$. Consider the following sequence:

$$C' \equiv (i^{M+1}, o^M, i^M, o^{M-1}, \dots, i^{m+1}, o^m).$$

Since we assume that $(i^1, o^1, \dots, i^M, o^M)$ satisfies **(C1)**, we have that $o^{m'} \notin \nu(i^{m'+1})$ and $o^{m'} \in \nu(i^{m'})$ for all $m' \in \{1, \dots, M+1\}$. Hence, C' is a cycle of ν . We now show that C' is a GCIC of ν for i^{M+1} at A . By **(C2)** above, $o^{m'} \in \Omega_{i^{m'+1}} \cup A_{i^{m'+1}}$ for all $m' \in \{1, \dots, M\}$, so that C' is individually rational at A . Next, note that **(C3)** and $i^{M+1} < t$ imply that C' does not affect any agent with higher priority than i^{M+1} . Finally, **(C3)**, $i^{M+1} < t$, and $o^M \in A_{i^{M+1}}$ imply the remaining requirements for C' to be a GCIC of ν for i^{M+1} at A . Since C' is a GCIC of ν for i^{M+1} at A , by Lemma 5, $\nu \notin \mathcal{M}^{i^{M+1}}(A)$. Since $i^{M+1} < t$, we have $\mathcal{M}^t(A) \subseteq \mathcal{M}^{i^{M+1}}(A)$ and thus $\nu \notin \mathcal{M}^t(A)$, which is a contradiction. Thus, $o^m \in A_{i^{M+1}}$.

Since m was arbitrary, **(C3)** and the just established fact imply that, for each $m' \leq M$ such that $i^{m'} = i^{M+1}$, $\{o^{m'-1}, o^{m'}\} \subseteq A_{i^{M+1}}$. Since $\nu, \mu \in \mathcal{M}^{t-1}(A) \subseteq \mathcal{M}^{i^{M+1}}(A)$, we have $|\nu(i^{M+1}) \cap A_{i^{M+1}}| = |\mu(i^{M+1}) \cap A_{i^{M+1}}|$. Since $o^M \in (\mu(i^{M+1}) \setminus \nu(i^{M+1})) \cap A_{i^{M+1}}$ and, for each $m' \in \{2, \dots, M-1\}$, $o^{m'-1} \in \mu(i^{m'}) \setminus \nu(i^{m'})$ and $o^{m'} \in \nu(i^{m'}) \setminus \mu(i^{m'})$, we have that

$$\begin{aligned} & |(\nu(i^{M+1}) \setminus \mu(i^{M+1})) \cap \{o^1, \dots, o^M\} \cap A_{i^{M+1}}| \\ & \quad < \\ & |(\mu(i^{M+1}) \setminus \nu(i^{M+1})) \cap \{o^1, \dots, o^M\} \cap A_{i^{M+1}}|. \end{aligned}$$

Thus, combining the last two observations, there exists $o^{M+1} \in [(\nu(i^{M+1}) \setminus \mu(i^{M+1})) \setminus \{o^1, \dots, o^M\}] \cap A_{i^{M+1}}$.

Case 2.2: $i^{M+1} < t$ and $o^M \in \Omega_{i^{M+1}} \setminus A_{i^{M+1}}$ We first show that $o^m \in \Omega_{i^{M+1}} \setminus A_{i^{M+1}}$. Otherwise, $o^m \in A_{i^{M+1}}$. Consider the following sequence:

$$(i^m, o^m, \dots, i^M, o^M)$$

Since $o^m \in A_{i_{M+1}}$ but $o^M \in \Omega_{i_{M+1}} \setminus A_{i_{M+1}}$, by the assumptions that we have made with regards to $(i^1, o^1, \dots, i^M, o^M)$, this sequence is a GCIC of μ for i^{M+1} at A . Hence, Lemma 5 implies that $\mu \notin \mathcal{M}^{i^{M+1}}(A)$. Since $\mu \in \mathcal{M}^t(A) \subseteq \mathcal{M}^{i^{M+1}}(A)$, we obtain a contradiction. Thus, $o^m \in \Omega_{i_{M+1}} \setminus A_{i_{M+1}}$.

Since m was arbitrary, (C3) and the just established fact imply that, for each $m' \leq M$ such that $i^{m'} = i^{M+1}$, $\{o^{m'-1}, o^{m'}\} \subseteq \Omega_{i_{M+1}}$. Since $\nu, \mu \in \mathcal{M}^{i^{M+1}}(A) \subseteq \mathcal{M}^{i^{M+1}}(A)$, we obtain $|\nu(i^{M+1}) \cap A_{i_{M+1}}| = |\mu(i^{M+1}) \cap A_{i_{M+1}}|$ and hence $|\nu(i^{M+1}) \cap (\Omega_{i_{M+1}} \setminus A_{i_{M+1}})| = |\mu(i^{M+1}) \cap (\Omega_{i_{M+1}} \setminus A_{i_{M+1}})|$. Since $o^M \in (\mu(i^{M+1}) \setminus \nu(i^{M+1})) \cap (\Omega_{i_{M+1}} \setminus A_{i_{M+1}})$ and, for each $m' \in \{2, \dots, M-1\}$, $o^{m'-1} \in \mu(i^{m'}) \setminus \nu(i^{m'})$ as well as $o^{m'} \in \nu(i^{m'}) \setminus \mu(i^{m'})$, we have that

$$\begin{aligned} & |(\nu(i^{M+1}) \setminus \mu(i^{M+1})) \cap \{o^1, \dots, o^M\} \cap (\Omega_{i_{M+1}} \setminus A_{i_{M+1}})| \\ & \qquad \qquad \qquad < \\ & |(\mu(i^{M+1}) \setminus \nu(i^{M+1})) \cap \{o^1, \dots, o^M\} \cap (\Omega_{i_{M+1}} \setminus A_{i_{M+1}})|. \end{aligned}$$

Thus, there is $o^{M+1} \in [(\nu(i^{M+1}) \setminus \mu(i^{M+1})) \setminus \{o^1, \dots, o^M\}] \cap (\Omega_{i_{M+1}} \setminus A_{i_{M+1}})$.

Case 2.3: $i^{M+1} > t$ Since ν and μ are both feasible, $|\mu(i^{M+1}) \setminus \nu(i^{M+1})| = |\nu(i^{M+1}) \setminus \mu(i^{M+1})|$. However, since $o^M \in \mu(i^{M+1}) \setminus \nu(i^{M+1})$ and for each $m \in \{2, \dots, M-1\}$, $o^{m-1} \in \mu(i^m) \setminus \nu(i^m)$ as well as $o^m \in \nu(o^m) \setminus \mu(o^m)$, we have that

$$|(\nu(i^{M+1}) \setminus \mu(i^{M+1})) \cap \{o^1, \dots, o^M\}| < |(\mu(i^{M+1}) \setminus \nu(i^{M+1})) \cap \{o^1, \dots, o^M\}|.$$

Thus, there is $o^{M+1} \in (\nu(i^{M+1}) \setminus \mu(i^{M+1})) \setminus \{o^1, \dots, o^M\}$.

Case 3: $i^{M+1} = t$ and $o^M \in A_{i_{M+1}}$ By (C4), $|(\nu(t) \setminus \mu(t)) \cap \{o^1, \dots, o^M\} \cap A_t| = |(\mu(t) \setminus \nu(t)) \cap \{o^1, \dots, o^M\} \cap A_t|$. Since $|\nu(t) \cap A_t| > |\mu(t) \cap A_t|$, there is $o^{M+1} \in [(\nu(t) \setminus \mu(t)) \setminus \{o^1, \dots, o^M\}] \cap A_t$.¹⁰

C Proofs from Section 4

Proof of Lemma 1. Let j be the highest priority agent for whom $K^j(\hat{A}) \neq K^j(A)$ and note that $K^i(A) \neq K^i(\hat{A})$ implies $j \leq i$.

We first show via induction on k that, for all $k \leq j$, $\mathcal{M}^{k-1}(A) \subseteq \mathcal{M}^{k-1}(\hat{A})$ and $o \in \mu(i)$ for each $\mu \in \mathcal{M}^{k-1}(\hat{A}) \setminus \mathcal{M}^{k-1}(A)$. For $k = 1$, both statements follow from $\hat{A}_i = A_i \cup \{o\}$ and $\hat{A}_l = A_l$ for all $l \neq i$. So consider some $k \in \{2, \dots, j\}$ and assume that both statements

¹⁰Recall that $i^{M+1} = t$ implies $o^M \in A_{i_{M+1}}$ (as otherwise there would have been no need to extend the sequence $i^1, o^1, \dots, i^M, o^M$) so Cases 1, 2, and 3 are exhaustive.

have been shown for all $k' < k$. We have that

$$\begin{aligned}
\mathcal{M}^{k-1}(A) &= \{\mu \in \mathcal{M}^{k-2}(A) : |\mu(k-1) \cap A_{k-1}| = K^{k-1}(A)\} \\
&\subseteq \{\mu \in \mathcal{M}^{k-2}(\hat{A}) : |\mu(k-1) \cap A_{k-1}| = K^{k-1}(A)\} \\
&= \{\mu \in \mathcal{M}^{k-2}(\hat{A}) : |\mu(k-1) \cap \hat{A}_{k-1}| = K^{k-1}(\hat{A})\} \\
&= \mathcal{M}^{k-1}(\hat{A})
\end{aligned}$$

Here, the subset relation follows from the inductive assumption that $\mathcal{M}^{k-2}(A) \subseteq \mathcal{M}^{k-2}(\hat{A})$ and the second equality follows from the definitions of j and \hat{A} since $k-1 < j \leq i$. In order to show the second part of the statement for k , fix an arbitrary $\mu \in \mathcal{M}^{k-1}(\hat{A})$ such that $o \notin \mu(i)$. By construction of $\mathcal{M}^{k-1}(\hat{A})$, we have $\mu \in \mathcal{M}^{k-2}(\hat{A})$. Since $o \notin \mu(i)$, the inductive assumption for $k-1$ implies $\mu \in \mathcal{M}^{k-2}(A)$. Since $k-1 < j \leq i$, we have that $|\mu(k-1) \cap A_{k-1}| = |\mu(k-1) \cap \hat{A}_{k-1}|$ and $K^{k-1}(A) = K^{k-1}(\hat{A})$. Combining the last two statements, we obtain $\mu \in \mathcal{M}^{k-1}(A)$.

We now use the just established statements for $k = j$ to complete the proof. Since $\mathcal{M}^{j-1}(A) \subseteq \mathcal{M}^{j-1}(\hat{A})$, $K^j(\hat{A}) \geq K^j(A)$ and the definition of j implies $K^j(\hat{A}) > K^j(A)$. Thus, $\mathcal{M}^j(\hat{A}) \subseteq \mathcal{M}^{j-1}(\hat{A}) \setminus \mathcal{M}^{j-1}(A)$. Hence, $o \in \mu(i)$ for all $\mu \in \mathcal{M}^j(\hat{A})$. Since $j \leq i$, we have that $\mathcal{M}^i(\hat{A}) \subseteq \mathcal{M}^j(\hat{A})$. Combining the last two observations we obtain that $o \in \mu(i)$ for all $\mu \in \mathcal{M}^i(\hat{A})$, which proves the desired statement. \square

Proof of Lemma 2. Since $\Omega_i \cup A_i = \Omega_i \cup \hat{A}_i$, $\mathcal{M}^0(\hat{A}) = \mathcal{M}^0(A)$. Since, for each $t < i$, $\hat{A}_t = A_t$, $\mathcal{M}^t(\hat{A}) = \mathcal{M}^t(A)$. Thus, $\mathcal{M}^{i-1}(\hat{A}) = \mathcal{M}^{i-1}(A)$. By definition of $K^i(A)$ and $\mathcal{M}^i(A)$, for each $\mu \in \mathcal{M}^i(A) \subseteq \mathcal{M}^{i-1}(\hat{A})$, $K^i(A) \geq |\mu(i) \cap A_i|$. \square

Proof of Lemma 3. If $K^i(A) = K^i(\hat{A})$, there is nothing left to show. So for the remainder of the proof, assume that $K^i(A) \neq K^i(\hat{A})$. We argue that $K^i(A) > K^i(\hat{A})$.

Let j^* be the highest priority agent for whom $K^{j^*}(\hat{A}) \neq K^{j^*}(A)$. Note that $K^i(A) \neq K^i(\hat{A})$ implies $j^* \leq i$.

We argue first that $K^{j^*}(A) > K^{j^*}(\hat{A})$. Since $K^j(\hat{A}) = K^j(A)$ for all $j < j^*$ and $\hat{A}_k \subseteq A_k$ for all $k \in I$, we have $\mathcal{M}^{j^*-1}(\hat{A}) \subseteq \mathcal{M}^{j^*-1}(A)$. The last subset relation implies $K^{j^*}(A) \geq K^{j^*}(\hat{A})$ and thus, given that we assumed $K^{j^*}(A) \neq K^{j^*}(\hat{A})$, we have $K^{j^*}(A) > K^{j^*}(\hat{A})$.

Given that $K^{j^*}(A) > K^{j^*}(\hat{A})$, the statement of Lemma 3 follows if $j^* = i$. Henceforth, we assume that $j^* < i$. We show that, for each $\mu \in \mathcal{M}^{i-1}(\hat{A})$, there is a CIC C^* of μ for j^* at A such that $\mu + C^* \in \mathcal{M}^{i-1}(A)$ and $|(\mu + C^*)(i) \cap A_i| \geq |\mu(i) \cap A_i|$. In particular, for any $\mu \in \mathcal{M}^{i-1}(\hat{A})$ there exists a $\mu' \in \mathcal{M}^{i-1}(A)$ such that $|\mu'(i) \cap A_i| \geq |\mu(i) \cap A_i|$. Hence, $K^i(A) \geq K^i(\hat{A})$ and thus, given that we have assumed $K^i(A) \neq K^i(\hat{A})$, $K^i(A) > K^i(\hat{A})$, which yields Lemma 3.

For the remainder of the proof of Lemma 3, we fix an arbitrary $\mu \in \mathcal{M}^{i-1}(\hat{A})$ and show that a CIC of μ for j^* at A with the desired properties exists. Since the proof is quite involved, we introduce eight claims.

Our first claim is that μ *complies with the promise to agents* $j' \leq j^* - 1$ in the IRP algorithm under A (i.e. satisfies $\mu \in \mathcal{M}^{j^*-1}(A)$) but not with the promise to j^* (i.e. $\mu \notin \mathcal{M}^{j^*}(A)$).

Claim 1. $\mu \in \mathcal{M}^{j^*-1}(A) \setminus \mathcal{M}^{j^*}(A)$

By Theorem 2, Claim 1 implies that there exists a CIC $C \equiv (i^1, o^1, \dots, i^M, o^M)$ of μ for j^* at A . The CIC C remains fixed throughout the remainder of the proof of Lemma 3. Our second claim is that i receives o in C .

Claim 2. $o \in (\mu + C)(i) \setminus \mu(i)$

If there is no CIC of $\mu + C$ for any $j < i$ at A . By Claim 1 and Theorem 2, we have that $\mu + C \in \mathcal{M}^{i-1}(A)$. By Claim 2, we have that $o \in (\mu + C)(i) \setminus \mu(i)$ and thus, given that $o \in A_i$, $|(\mu + C)(i) \cap A_i| \geq |\mu(i) \cap A_i|$. Hence, C is a CIC with the desired properties and we are done.

Now, we consider the case where there is an agent $\hat{j}^* < i$ and a CIC $\hat{C} \equiv (j^1, p^1, \dots, j^L, p^L)$ of $\mu + C$ for \hat{j}^* at A . The CIC \hat{C} remains fixed throughout the remainder of the proof of Lemma 3. We assume without loss of generality that there is no agent $j' < \hat{j}^*$ for whom there is a CIC of $\mu + C$ at A . Our next claim lists two important initial observations about \hat{j}^* , C , and \hat{C} .

Claim 3. $\hat{j}^* \geq j^*$ and $\{\mu + C, (\mu + C) + \hat{C}\} \subseteq \mathcal{M}^{\hat{j}^*-1}(A)$

In the following, we use C and \hat{C} to show that there exists a third CIC \tilde{C} of μ for j^* at A that satisfies the following properties:

$$\mu + \tilde{C} \in \mathcal{M}^{\hat{j}^*-1}(A) \quad (\text{P1})$$

$$|(\mu + \tilde{C})(\hat{j}^*) \cap A_{\hat{j}^*}| \geq |((\mu + C) + \hat{C})(\hat{j}^*) \cap A_{\hat{j}^*}| \quad (\text{P2})$$

$$|(\mu + \tilde{C})(i) \cap A_i| \geq |\mu(i) \cap A_i| \quad (\text{P3})$$

Repeated application of this argument allows us to infer that there exists a CIC C^* of μ for j^* at A that does not decrease the welfare of i and that satisfies $\mu + C^* \in \mathcal{M}^{i-1}(A)$.¹¹ As

¹¹To see this, note first that if $\mu + \tilde{C} \notin \mathcal{M}^{i-1}(A)$, then Theorem 2 and $\mu + \tilde{C} \in \mathcal{M}^{\hat{j}^*-1}(A)$ imply that there exists a CIC \tilde{C}' of $\mu + \tilde{C}$ for some $j' \geq \hat{j}^*$. We can now apply our earlier finding to \tilde{C} and \tilde{C}' to get yet another CIC that satisfies the three properties listed above. Finally, note if $j' = \hat{j}^*$, then we have that $|((\mu + \tilde{C}) + \tilde{C}')(\hat{j}^*) \cap A_{\hat{j}^*}| > |(\mu + \tilde{C})(\hat{j}^*) \cap A_{\hat{j}^*}|$ since $|(\mu + \tilde{C})(\hat{j}^*) \cap A_{\hat{j}^*}| \geq |((\mu + C) + \hat{C})(\hat{j}^*) \cap A_{\hat{j}^*}| \geq |\mu(\hat{j}^*) \cap A_{\hat{j}^*}|$ (the second inequality follows since \hat{C} is a CIC for \hat{j}^*) and since \tilde{C}' is a CIC for \hat{j}^* . Since the set of agents is finite and since the maximum number of desirable objects an agent can obtain is finite, we eventually find a CIC with the desired properties.

mentioned above, the existence of such a CIC proves Lemma 3.

Note that since $(\mu + C) + \hat{C} \in \mathcal{M}^{\hat{j}^*-1}(A)$, a CIC \tilde{C} satisfies (P1) - (P3) if and only if

1. $\mu + \tilde{C}$ makes every agent $j' \leq \hat{j}^*$ at least as well off as $(\mu + C) + \hat{C}$, i.e. $|(\mu + \tilde{C})(j') \cap A_{j'}| \geq |((\mu + C) + \hat{C})(j') \cap A_{j'}|$ for all $j' \leq \hat{j}^*$, and
2. \tilde{C} does not decrease the welfare of i , i.e. $|(\mu + \tilde{C})(i) \cap A_i| \geq |\mu(i) \cap A_i|$.

In our arguments below, we always show that \tilde{C} satisfies these two properties.

We start by introducing some basic notation. Given two integers $m, m' \in \{1, \dots, M\}$, we use the notation

- $\{m, \dots [M, 1] \dots, m'\}$ to represent $\{m, \dots, m'\}$ if $m \leq m'$ and $\{1, \dots, m'\} \cup \{m, \dots, M\}$ if $m > m'$,
- $(i^m, o^m, \dots [o^M, i^1] \dots, i^{m'}, o^{m'})$ to represent $(i^m, o^m, \dots, i^{m'}, o^{m'})$ if $m \leq m'$ and $(i^m, o^m, \dots, i^M, o^M, i^1, o^1, \dots, i^{m'}, o^{m'})$ if $m > m'$,
- $(i^m, \dots [i^M, i^1] \dots, i^{m'})$ to represent $(i^m, \dots, i^{m'})$ if $m \leq m'$ and $(i^m, \dots, i^M, i^1, \dots, i^{m'})$ if $m > m'$, and
- $(o^m, \dots [o^M, o^1] \dots, o^{m'})$ to represent $(o^m, \dots, o^{m'})$ if $m \leq m'$ and $(o^m, \dots, o^M, o^1, \dots, o^{m'})$ if $m > m'$.

Analogously, given two integers $l, l' \in \{1, \dots, L\}$, we use the notations $\{l, \dots [L, 1] \dots, l'\}$, $(j^n, p^l, \dots [p^L, j^1] \dots, j', p^{l'})$, $(j^l, \dots [j^L, j^1] \dots, j^{l'})$, and $(p^l, \dots [p^L, p^1] \dots, p^{l'})$.

Next, we derive two basic properties of C and \hat{C} . First, we establish that C and \hat{C} “intersect” in terms of objects.

Claim 4. $O(C) \cap O(\hat{C}) \neq \emptyset$

Next, we argue that C and \hat{C} treat agents with higher priority than j^* “consistently”.

Claim 5. For each pair m and l such that $i^m = j^l \equiv \underline{j}$ and $\underline{j} < j^*$, either

$$\{o^{m-1}, o^m, p^{l-1}, p^l\} \subseteq A_{\underline{j}} \text{ or } \{o^{m-1}, o^m, p^{l-1}, p^l\} \subseteq \Omega_{\underline{j}} \setminus A_{\underline{j}}.$$

We now introduce some further notation. First, let \underline{l}^* be the smallest integer such that either $j^{\underline{l}^*} \in I(C) \setminus \{\hat{j}^*\}$ or $p^{\underline{l}^*} \in O(C)$. Claim 4 ensures that such \underline{l}^* exists. We now use \underline{l}^* to define a specific point on the cycle C . In doing so, we distinguish two cases:

1. If $j^{\underline{l}^*} \notin I(C) \setminus \{\hat{j}^*\}$ and $p^{\underline{l}^*} \in O(C)$, then let $\underline{l} = \underline{l}^*$ and let $m(\underline{l})$ be such that $p^{\underline{l}} = o^{m(\underline{l})-1}$ (where $o^0 \equiv o^M$).

2. If $j^{\underline{l}^*} \in I(C) \setminus \{\hat{j}^*\}$, then let $\underline{l} = \underline{l}^* - 1$ and let $m(\underline{l})$ be such that $j^{\underline{l}^*} = i^{m(\underline{l})}$.¹²

Note that in both two cases, our definition of \underline{l} ensures that $\underline{l} \geq 1$, $j^{\underline{l}} \notin I(C) \setminus \{\hat{j}^*\}$, and $p^{\underline{l}} \in \mu(i^{m(\underline{l})})$. To understand the significance of $m(\underline{l})$, note first that the definition of \underline{l} implies that $\{j^1, p^1, \dots, j^{\underline{l}}, p^{\underline{l}}\} \cap \{i^1, o^1, \dots, i^M, o^M\} \subseteq \{j^1, p^{\underline{l}}\}$. Now consider the following sequence of agent-object pairs:

$$\underline{C} \equiv (j^1, p^1, \dots, j^{\underline{l}}, p^{\underline{l}}, i^{m(\underline{l})}, o^{m(\underline{l})}, \dots, i^M, o^M).$$

In \underline{C} , each agent points to an object that she does not get at μ and each object, except possibly o^M , points to the agent who gets it at μ . Furthermore, except possibly j^1 , all agents and objects in \underline{C} are distinct from each other. These facts about \underline{C} are useful for our construction of CICs in the remainder of the proof.

Second, let \bar{l} be the largest integer such that either $j^{\bar{l}} \in I(C) \setminus \{\hat{j}^*\}$ or $p^{\bar{l}} \in O(C)$. Again, Claim 4 ensures that such \bar{l} exists. As is the case for \underline{l}^* , we use \bar{l} to define a specific point of the cycle C . Before doing that, we discuss the case of $p^{\bar{l}} \in O(C)$ in more detail by means of the following claim.

Claim 6. *If $p^{\bar{l}} \in O(C)$, then $\bar{l} = L$.*

Now consider the case where $p^{\bar{l}} \notin O(C)$ and $j^{\bar{l}} \in I(C) \setminus \{\hat{j}^*\}$. Let $m(\bar{l})$ be such that $i^{m(\bar{l})+1} = j^{\bar{l}}$. Since C is a CIC of μ , $j^{\bar{l}} = i^{m(\bar{l})+1}$ implies that $o^{m(\bar{l})} \in \mu(j^{\bar{l}})$. By definition of \bar{l} , we have that $\{i^1, o^1, \dots, i^M, o^M\} \cap \{j^{\bar{l}}, p^{\bar{l}}, \dots, j^L, p^L\} = \{j^{\bar{l}}\}$. Now let $m' \in \{1, \dots, M\}$ be an arbitrary integer distinct from $m(\bar{l}) + 1$ and consider the following sequence of agent-object pairs:

$$\bar{C} \equiv (i^{m'}, o^{m'}, \dots, [o^M, i^1] \dots, i^{m(\bar{l})}, o^{m(\bar{l})}, j^{\bar{l}}, p^{\bar{l}}, \dots, j^L, p^L).$$

In \bar{C} , each agent points to an object that she does not get at μ and each object, except possibly p^L , points to the agent who gets it at μ . Furthermore, all agents and objects in \bar{C} are distinct from each other. Again, these facts are useful for our construction of CICs in the remainder of the proof.

Before proceeding, we now complete the proof of Lemma 3 for the case of $I(C) \cap I(\hat{C}) = \{j^*\}$. Let l be such that $j^l = j^*$. Note that since $I(C) \cap I(\hat{C}) = \{j^*\}$, Claim 4 implies $O(C) \cap O(\hat{C}) = \{o^1\}$ as well as $p^{l-1} = o^1$. If $j^* = \hat{j}^*$, we arrive at a contradiction since $p^L \in \Omega_{\hat{j}^*} \setminus A_{\hat{j}^*}$ (as \hat{C} is a CIC for \hat{j}^*) and $o^1 \in A_{j^*}$ (as C is a CIC for j^*) imply $p^L \neq o^1$. Thus, $j^* \neq \hat{j}^*$. So, by Claim 3, $j^* < \hat{j}^*$. Since \hat{C} is a CIC for \hat{j}^* , we obtain that $p^l \in A_{j^*}$.

¹²Note that $\underline{l}^* \geq 2$ in this case.

Given that $I(C) \cap I(\hat{C}) = \{j^*\}$ and $O(C) \cap O(\hat{C}) = \{o^1\}$,

$$\tilde{C} \equiv (j^l, p^l, \dots, [p^L, j^1], \dots, j^{l-1}, p^{l-1}, i^2, o^2, \dots, i^M, o^M)$$

is a CIC of μ for j^* at A such that $\mu + \tilde{C}$ makes all agents exactly as well off as $(\mu + C) + \hat{C}$. This completes the proof of Lemma 3 in case $I(C) \cap I(\hat{C}) = \{j^*\}$.

Henceforth, we assume that $I(C) \cap I(\hat{C}) \neq \{j^*\}$. Claim 4 then implies $(I(C) \cap I(\hat{C})) \setminus \{j^*\} \neq \emptyset$.

Next, we introduce some further notation for the remainder of the proof in case $I(C) \cap I(\hat{C}) \neq \{j^*\}$. Let \underline{m} be the smallest integer m such that $i^m \in I(\hat{C}) \setminus \{j^*\}$, i.e. $i^{\underline{m}} \in I(\hat{C}) \setminus \{j^*\}$ and for each $m \in \{1, \dots, \underline{m} - 1\}$, $i^m \notin I(\hat{C}) \setminus \{j^*\}$.¹³ Similarly, let \overline{m} be the largest integer m' such that $i^{m'} \in I(\hat{C}) \setminus \{j^*\}$, i.e. $i^{\overline{m}} \in I(\hat{C}) \setminus \{j^*\}$ and for each $m \in \{\overline{m} + 1, \dots, M\}$, $i^m \notin I(\hat{C}) \setminus \{j^*\}$. Note that it is possible that $\overline{m} = \underline{m}$ and that in this case, we have $(I(C) \cap I(\hat{C})) \setminus \{j^*\} = \{i^{\overline{m}}\}$, $o^{\overline{m}} \in O(\hat{C})$, and $O(C) \cap O(\hat{C}) \subseteq \{o^1, o^{\overline{m}}\}$. Finally, let m^* be such that $i^{m^*} = i$ and $o^{m^*} = o$. Note that the existence of such an m^* follows since $o \in (\mu + C)(i) \setminus \mu(i)$ by Claim 2.

Our next claim is that objects and agents that show up in both C and \hat{C} either all appear “before”, or all appear “after” agent i in C , starting from $i^1 (= j^*)$.

Claim 7. $o \notin O(\hat{C})$ and if $\underline{m} \leq m^*$, then $\overline{m} \leq m^*$.

Our final claim lays the foundation for splicing together parts of C and \hat{C} to form new CICs. As we explain below, the claim will allow us to cut and paste parts of C and \hat{C} together to construct a CIC that makes agents $j' \in \{j^*, \dots, \hat{j}^* - 1\}$ at least as well off as $(\mu + C) + \hat{C}$.

Claim 8. Assume that $\underline{m} < \overline{m}$.

1. Let $\check{m} \in \{\underline{m} + 1, \dots, \overline{m} - 1\}$ be such that $i^{\check{m}} < \hat{j}^*$.

(i) The cycle C does not affect $i^{\check{m}}$.

(ii) If there exists \check{l} such that $j^{\check{l}} = i^{\check{m}}$, then either $\{o^{\check{m}-1}, o^{\check{m}}, p^{\check{l}-1}, p^{\check{l}}\} \subseteq A_{i^{\check{m}}}$ or $\{o^{\check{m}-1}, o^{\check{m}}, p^{\check{l}-1}, p^{\check{l}}\} \subseteq \Omega_{i^{\check{m}}} \setminus A_{i^{\check{m}}}$.

2. Let $\check{m} \leq \underline{m}$ be such that $i^{\check{m}} < \hat{j}^*$.

(i) If C affects $i^{\check{m}}$, then either $j^* \notin I(\hat{C})$ or there exists l^* such that $j^{l^*} = j^*$ and $\{p^{l^*-1}, p^{l^*}\} \subseteq \Omega_{j^*} \setminus A_{j^*}$.

(ii) If $\check{m} = \underline{m}$, then $o^{\check{m}} \in A_{i^{\check{m}}}$ if and only if there exists a \check{l} such that $j^{\check{l}} = i^{\check{m}}$ and $\{p^{\check{l}-1}, p^{\check{l}}\} \subseteq A_{i^{\check{m}}}$.

¹³Note that the existence of such an integer follows from $(I(C) \cap I(\hat{C})) \setminus \{j^*\} \neq \emptyset$.

3. Let $\tilde{m} \geq \bar{m}$ be such that $i^{\tilde{m}} < \hat{j}^*$.

(i) If C affects $i^{\tilde{m}}$, then either $j^* \notin I(\hat{C})$ or there exists l^* such that $j^{l^*} = j^*$ and $\{p^{l^*-1}, p^{l^*}\} \subseteq A_{j^*}$.

(ii) If $\tilde{m} = \bar{m}$, then $o^{\bar{m}-1} \in A_{i^{\bar{m}}}$ if and only if there exists a \tilde{l} such that $j^{\tilde{l}} = i^{\bar{m}}$ and $\{p^{\tilde{l}-1}, p^{\tilde{l}}\} \subseteq A_{i^{\bar{m}}}$.

4. If \tilde{m} is such that $i^{\tilde{m}} = \hat{j}^*$, then C does not increase the welfare of $i^{\tilde{m}}$.

We now use the above claims to show that either there is a CIC \tilde{C} of μ for j^* at A that satisfies (P1) - (P3), or there is a CIC of μ for some agent $j' < i$ at \hat{A} (contradicting $\mu \in \mathcal{M}^{i-1}(\hat{A})$), or there is a CIC of $\mu + C$ for some agent $j' < \hat{j}^*$ (contradicting our choice of \hat{C}). Since we only work with the reassignments implied by C and \hat{C} , Claim 5 implies that the CICs we construct cannot affect any agent $j' < j^*$. We use this observation repeatedly throughout the remainder of our proof. To structure our arguments, we distinguish 5 cases.

Case 1: $\underline{m} = \bar{m}$. As argued above, $I(C) \cap I(\hat{C}) \subseteq \{j^*, i^{\underline{m}}\}$ and $O(C) \cap O(\hat{C}) \subseteq \{o^1, o^{\underline{m}}\}$.

Assume first that $o^{\underline{m}} \in O(\hat{C})$. Let \tilde{l} be such that $j^{\tilde{l}} = i^{\underline{m}}$ and note that $o^{\underline{m}} = p^{\tilde{l}-1}$ in the case we consider here. In particular, it is not possible that C and \hat{C} both increase the welfare of $i^{\underline{m}}$. If $j^* \notin I(\hat{C})$, then

$$\tilde{C} \equiv (i^1, o^1, \dots, i^{\underline{m}-1}, o^{\underline{m}-1}, j^{\tilde{l}}, p^{\tilde{l}}, \dots [p^L, j^1] \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1}, i^{\underline{m}+1}, o^{\underline{m}+1}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at A that satisfies (P1) - (P3): Since $I(C) \cap I(\hat{C}) \subseteq \{j^*, i^{\underline{m}}\}$ and $O(C) \cap O(\hat{C}) \subseteq \{o^1, o^{\underline{m}}\}$ and since C and \hat{C} are CICs for j^* and \hat{j}^* respectively, \tilde{C} is a CIC of μ for j^* at A . If $j^{\tilde{l}} < \hat{j}^*$, then since $o^{\underline{m}} = p^{\tilde{l}-1} \in A_{j^{\tilde{l}}}$ if and only if $p^{\tilde{l}} \in A_{j^{\tilde{l}}}$ given that \hat{C} is a CIC for $\hat{j}^* > j^{\tilde{l}}$, we conclude that $\mu + \tilde{C}$ makes $j^{\tilde{l}}$ at least as well off as $(\mu + C) + \hat{C}$. If $j^{\tilde{l}} = \hat{j}^*$ (and thus $\tilde{l} = 1$), then since $o^{\underline{m}} = p^{\tilde{l}-1} = p^L \in \Omega_{\hat{j}^*} \setminus A_{\hat{j}^*}$ and $p^1 \in A_{j^{\tilde{l}}}$, $j^{\tilde{l}}$ is at least well off under $\mu + \tilde{C}$ as under $(\mu + C) + \hat{C}$. Finally, $m^* \neq \underline{m}$ since otherwise \tilde{C} is a CIC of μ for j^* at \hat{A} . Hence, \tilde{C} weakly increases the welfare of i . Combining the previous arguments, we find that \tilde{C} is a CIC with the desired properties.

Now suppose that $j^* \in I(\hat{C})$ so there exists an $l^* > 1$ such that $j^{l^*} = j^*$.

If $p^{l^*-1} \in \Omega_{j^*} \setminus A_{j^*}$, then $o^1 \notin O(\hat{C})$. We claim that

$$\tilde{C}' \equiv (i^1, o^1, \dots, i^{\underline{m}-1}, o^{\underline{m}-1}, j^{\tilde{l}}, p^{\tilde{l}}, \dots [p^L, j^1] \dots, j^{l^*-1}, p^{l^*-1})$$

is a CIC of μ for j^* at A that satisfies (P1) - (P3): Since $I(C) \cap I(\hat{C}) \subseteq \{j^*, i^{\underline{m}}\}$ and $O(C) \cap O(\hat{C}) \subseteq \{o^1, o^{\underline{m}}\}$ and since C and \hat{C} are CICs for j^* and \hat{j}^* respectively, \tilde{C}' is a

CIC of μ for j^* at A . Since $\mu \in \mathcal{M}^{i-1}(\hat{A})$, we have $m^* \leq \underline{m} - 1$. As before, \tilde{C}' weakly increases the welfare of \hat{j}^* if $j^{\tilde{l}} = \hat{j}^*$. If $j^{\tilde{l}} \neq \hat{j}^*$, note first that a simple case distinction shows that there cannot exist a $\tilde{m} > \underline{m}$ such that $i^{\tilde{m}} < \hat{j}^*$ and such that C affects $i^{\tilde{m}}$.¹⁴ Hence, given that $m^* \leq \underline{m}$, $l^* - 1 \geq \tilde{l} > 1$ implies that

$$(j^1, p^1, \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1}, i^{\underline{m}+1}, o^{\underline{m}+1}, \dots, i^M, o^M)$$

is a CIC of μ for \hat{j}^* at \hat{A} . If $l^* - 1 < \tilde{l}$, \tilde{C}' weakly increases the welfare of \hat{j}^* compared to $(\mu + C) + \hat{C}$. Finally, as in the case of $j^* \notin I(\hat{C})$, one can show that \tilde{C}' cannot make $j^{\tilde{l}}$ worse off than $(\mu + C) + \hat{C}$ if $j^{\tilde{l}} < \hat{j}^*$.

If $p^{l^*-1} \in A_{j^*}$, then by analogous arguments

$$(j^{l^*}, p^{l^*}, \dots [p^L, j^1] \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1}, i^{\underline{m}+1}, o^{\underline{m}+1}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at A that satisfies (P1) - (P3). We omit the details.

Finally, suppose $O(C) \cap O(\hat{C}) = \{o^1\}$. Let l^* be such that $j^{l^*} = j^*$ and note that $p^{l^*-1} = o^1$ and $p^{l^*} \in A_{j^*}$. By analogs of the above arguments

$$\tilde{C}'' \equiv (j^{l^*}, p^{l^*}, \dots [p^L, j^1] \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1}, i^{\underline{m}}, o^{\underline{m}}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at A that satisfies (P1) - (P3).¹⁵

Case 2: $\underline{m} < \overline{m}$ and C decreases the welfare of \hat{j}^* . Let \tilde{m} be such that $i^{\tilde{m}} = \hat{j}^*$. The premise of Case 2 imply that there exist m_1, m_2, l_1, l_2 such that the following conditions are satisfied:

$$1. \underline{m} \leq m_1 \leq \tilde{m} \leq m_2 \leq \overline{m}$$

¹⁴Suppose, the contrary, that such an $i^{\tilde{m}}$ exists. Assume without loss of generality that there is no $m' \in \{\underline{m} + 1, \dots, M\}$ such that C affects $i^{m'}$ and such that $i^{m'} < i^{\tilde{m}}$. If C decreases the welfare of $i^{\tilde{m}}$, then

$$(i^1, o^M, i^M, o^{M-1}, \dots, i^{\underline{m}+1}, o^{\underline{m}}, j^{\tilde{l}}, p^{\tilde{l}}, \dots [p^L, j^1] \dots, j^{l^*-1}, p^{l^*-1})$$

is a CIC of $\mu + C$ for $i^{\tilde{m}}$ at A , which contradicts our choice of \hat{C} . If C increases the welfare of $i^{\tilde{m}}$, then

$$(i^{\underline{m}+1}, o^{\underline{m}+1}, \dots, i^M, o^M, j^{l^*}, p^{l^*}, \dots [p^L, j^1] \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1})$$

is a CIC of μ for $i^{\tilde{m}}$ at \hat{A} given that $m^* \leq \underline{m}$.

¹⁵One difference to the case is that when $o^{\underline{m}} \notin O(\hat{C})$, it is possible for C and \hat{C} to both increase the welfare of $i^{\underline{m}}$ and $i^{\underline{m}} < \hat{j}^*$. However, since \tilde{C}'' is a CIC of μ for j^* at A , we have $m^* \geq \underline{m}$ and

$$(j^{\tilde{l}}, p^{\tilde{l}}, \dots [p^L, j^1] \dots, j^{l^*-1}, p^{l^*-1}, i^2, o^2, \dots, i^{\underline{m}-1}, o^{\underline{m}-1})$$

is a CIC of μ for $i^{\underline{m}}$ at \hat{A} given that $p^{\tilde{l}} \in A_{j^i}$ and $o^{\underline{m}-1} \in \Omega_{j^i} \setminus A_{j^i}$.

2. Either $m_1 < \check{m}$ or $\check{m} < m_2$.
3. $j^{l_1} = i^{m_1}$ and $j^{l_2} = i^{m_2}$
4. $\{j^{l_1+1}, \dots, [j^L, j^1], \dots, j^{l_2-1}\} \cap I(C) \subseteq \{j^*, \hat{j}^*\}$

Assume first that $\{j^{l_1+1}, \dots, [j^L, j^1], \dots, j^{l_2-1}\} \cap I(C) = \emptyset$. If $p^{l_2-1} \notin O(C)$, then

$$\tilde{C} \equiv (i^1, o^1, \dots, i^{m_1-1}, o^{m_1-1}, j^{l_1}, p^{l_1}, \dots, [p^L, j^1], \dots, j^{l_2-1}, p^{l_2-1}, i^{m_2}, o^{m_2}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at A that satisfies **(P1)** - **(P3)**.¹⁶ If $p^{l_2-1} \in O(C)$, then $p^{l_2-1} = o^{m_2}$, in which case

$$\tilde{C}' \equiv (i^1, o^1, \dots, i^{m_1-1}, o^{m_1-1}, j^{l_1}, p^{l_1}, \dots, [p^L, j^1], \dots, j^{l_2-1}, p^{l_2-1}, i^{m_2+1}, o^{m_2+1}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at A that satisfies **(P1)** - **(P3)**.¹⁷

Next, assume that there exists $l^* \in \{l_1 + 1, \dots, [L, 1], \dots, l_2 - 1\}$ such that $j^{l^*} = j^*$. If $\{p^{l^*-1}, p^{l^*}\} \subseteq A_{j^*}$, then either $p^{l_2-1} \notin O(C)$ or $p^{l_2-1} = o^{m_2}$. In the former case

$$(j^{l^*}, p^{l^*}, \dots, [p^L, j^1], \dots, j^{l_2-1}, p^{l_2-1}, i^{m_2}, o^{m_2}, \dots, i^M, o^M)$$

and in the latter case

$$(j^{l^*}, p^{l^*}, \dots, [p^L, j^1], \dots, j^{l_2-1}, p^{l_2-1}, i^{m_2+1}, o^{m_2+1}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at A that satisfies **(P1)** - **(P3)**.¹⁸ If $\{p^{l^*-1}, p^{l^*}\} \subseteq \Omega_{j^*} \setminus A_{j^*}$, then

$$(i^1, o^1, \dots, i^{m_1-1}, o^{m_1-1}, j^{l_1}, p^{l_1}, \dots, [p^L, j^1], \dots, j^{l^*-1}, p^{l^*-1})$$

¹⁶To see this, note first that since $\underline{m} \leq m_1 \leq m_2 \leq \bar{m}$ the first part of Claim 8 implies that there does not exist $m' \in \{m_1 + 1, \dots, m_2 - 1\}$ such that $i^{m'} < \hat{j}^*$ and such that C affects $i^{m'}$. Next, if $m_1 = \underline{m}$ ($m_2 = \bar{m}$) the second (third) part of Claim 8 implies that \tilde{C} makes i^{m_1} (i^{m_2}) at least as well off as $(\mu + C) + \tilde{C}$ if $i^{m_1} < \hat{j}^*$ ($i^{m_2} < \hat{j}^*$). For agent \hat{j}^* , the definitions of m_1, m_2, l_1, l_2 and the assumptions of Case 2 immediately imply that \tilde{C} weakly increases the welfare of \hat{j}^* . Finally, $\mu \in \mathcal{M}^{i-1}(\hat{A})$ implies that $m^* \in \{1, \dots, m_1 - 1, m_2, \dots, M\}$: otherwise \tilde{C} is a CIC of μ for j^* at \hat{A} . So \tilde{C} weakly increases the welfare of i . Analogous reasoning applies to all CICs that we construct in the remainder of the proof.

¹⁷Note that if $p^{l_2-1} = o^{m_2}$ and $i^{m_2} < \hat{j}^*$, then the third part of Claim 8 implies that C cannot increase the welfare of i^{m_2} if $m_2 = \bar{m}$ since $o^{m_2-1} \in \Omega_{i^{m_2}} \setminus A_{i^{m_2}}$ implies $\{p^{l_2-1}, p^{l_2}\} \subseteq \Omega_{i^{m_2}} \setminus A_{i^{m_2}}$ (and thus $o^{m_2} \in \Omega_{i^{m_2}} \setminus A_{i^{m_2}}$). Similar reasoning applies to all CICs that we construct in which we “skip” agents $i^{\underline{m}}$ or $i^{\bar{m}}$.

¹⁸Since j^* exchanges two desirable objects in the case we consider here, the first and second parts of Claim 8 implies that there cannot be an $m \in \{2, \dots, \bar{m} - 1\}$ such that $i^m < \hat{j}^*$ and such that C affects i^m . Analogous reasoning applies to all CICs that we construct for cases where j^* exchanges two desirable objects in \hat{C} .

is a CIC of μ for j^* at A that satisfies (P1) - (P3).¹⁹

Finally, consider the case where $\{j^{l_1+1}, \dots, [j^L, j^1], \dots, j^{l_2-1}\} \cap I(C) = \{\hat{j}^*\}$. Since all agents in \hat{C} are distinct, $m_1 < \check{m} < m_2$ in this case. If $p^L \notin O(C)$, then

$$(i^1, o^1, \dots, i^{m_1-1}, o^{m_1-1}, j^{l_1}, p^{l_1}, \dots, j^L, p^L, i^{\check{m}}, o^{\check{m}}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at A that satisfies (P1) - (P3). If $p^L = o^{\check{m}}$, then

$$(i^1, o^1, \dots, i^{m_1-1}, o^{m_1-1}, j^{l_1}, p^{l_1}, \dots, j^L, p^L, i^{\check{m}+1}, o^{\check{m}+1}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at A that satisfies (P1) - (P3).

Case 3: $\underline{m} < \bar{m}$, C does not affect \hat{j}^* , $p^L \notin O(C)$, and $j^{\bar{l}} \neq j^*$. By definition of $m(\bar{l})$ and the premise of this case, $m(\bar{l}) < M$ and $j^{\bar{l}} = o^{m(\bar{l})+1}$. Hence, $m(\bar{l}) < \bar{m}$. We will use this observation repeatedly throughout the proof in Case 3.

We first assume that $m(\underline{l}) \leq m(\bar{l})$.

We start by showing $m(\underline{l}) < \underline{m}$. Assume to the contrary that $m(\underline{l}) \geq \underline{m}$. Claim 7 and $m(\bar{l}) < \bar{m}$ implies that $m^* \notin \{m(\underline{l}), \dots, m(\bar{l})\}$. If $\hat{j}^* \notin \{i^{m(\underline{l})}, \dots, i^{m(\bar{l})}\}$, then

$$\tilde{C} \equiv (j^1, p^1, \dots, j^{\underline{l}}, p^{\underline{l}}, i^{m(\underline{l})}, o^{m(\underline{l})}, \dots, i^{m(\bar{l})}, o^{m(\bar{l})}, j^{\bar{l}}, p^{\bar{l}}, \dots, j^L, p^L)$$

is a CIC of μ for \hat{j}^* at \hat{A} .²⁰ If there exists a $\check{m} \in \{m(\underline{l}), \dots, m(\bar{l})\}$ such that $i^{\check{m}} = \hat{j}^*$ and $o^{\check{m}-1} \in \Omega_{\hat{j}^*} \setminus A_{\hat{j}^*}$, then

$$(j^1, p^1, \dots, j^{\underline{l}}, p^{\underline{l}}, i^{m(\underline{l})}, o^{m(\underline{l})}, \dots, i^{\check{m}-1}, o^{\check{m}-1})$$

is a CIC of μ for \hat{j}^* at \hat{A} . Finally, if there exists a $\check{m} \in \{m(\underline{l}), \dots, m(\bar{l})\}$ such that $i^{\check{m}} = \hat{j}^*$ and $o^{\check{m}-1} \in A_{\hat{j}^*}$, then, given that C does not affect \hat{j}^* , we have $o^{\check{m}} \in A_{\hat{j}^*}$ as well. Since $m^* \notin \{m(\underline{l}), \dots, m(\bar{l})\}$,

$$(i^{\check{m}}, o^{\check{m}}, \dots, i^{m(\bar{l})}, o^{m(\bar{l})}, j^{\bar{l}}, p^{\bar{l}}, \dots, j^L, p^L)$$

¹⁹Since j^* exchanges two undesirable objects in the case we consider here, the first and third parts of Claim 8 implies that there cannot be an $m \in \{\underline{m} + 1, \dots, M\}$ such that $i^m < \hat{j}^*$ and such that C affects i^m . Analogous reasoning applies to all CICs that we construct for cases where j^* exchanges two undesirable objects in \hat{C} .

²⁰By the first part of Claim 8, there does not exist $m \in \{\underline{m} + 1, \dots, \bar{m} - 1\}$ such that $i^m < \hat{j}^*$ and such that \tilde{C} affects i^m . If $m(\underline{l}) = \underline{m}$, then the second part of Claim 8 implies that \tilde{C} cannot affect $i^{m(\underline{l})}$ if $i^{m(\underline{l})} < \hat{j}^*$. Analogous reasoning applies to other cases where we construct CICs for \hat{j}^* .

is a CIC of μ for \hat{j}^* at \hat{A} . Since $m(\underline{l}) \geq \underline{m}$ necessarily leads to a contradiction, we have $m(\underline{l}) < \underline{m}$. Thus, we have established that $m(\underline{l}) < \underline{m}$.

Next, note that $m(\underline{l}) < \underline{m}$ implies that either $p^{\underline{l}} \in \mu(j^*)$ or $p^{\underline{l}} = o^1$.²¹

Assume first that $p^{\underline{l}} \in \mu(j^*) \cap (\Omega_{j^*} \setminus A_{j^*})$. If $\hat{j}^* \notin \{i^2, \dots, i^{m(\bar{l})}\}$, then

$$(i^1, o^1, \dots, i^{m(\bar{l})}, o^{m(\bar{l})}, j^{\bar{l}}, p^{\bar{l}}, \dots, [p^{\underline{l}}, j^1] \dots, j^{\underline{l}}, p^{\underline{l}}).$$

is a CIC of μ for j^* at A that satisfies **(P1)** - **(P3)**. If there exists a $\check{m} \in \{2, \dots, m(\bar{l})\}$ such that $i^{\check{m}} = \hat{j}^*$ and $\{o^{\check{m}-1}, o^{\check{m}}\} \subseteq \Omega_{j^*} \setminus A_{j^*}$, then

$$(i^1, o^1, \dots, i^{\check{m}-1}, o^{\check{m}-1}, j^1, p^1, \dots, j^{\underline{l}}, p^{\underline{l}})$$

is a CIC of μ for j^* at A that satisfies **(P1)** - **(P3)**. If there exists a $\check{m} \in \{2, \dots, m(\bar{l})\}$ such that $i^{\check{m}} = \hat{j}^*$ and $\{o^{\check{m}-1}, o^{\check{m}}\} \subseteq A_{j^*}$, note first that $m^* < \check{m}$ since otherwise either

$$(i^1, o^1, \dots, i^{\check{m}-1}, o^{\check{m}-1}, j^1, p^1, \dots, j^{\underline{l}}, p^{\underline{l}})$$

(if $o^{\check{m}-1} \neq p^1$) or

$$(i^1, o^1, \dots, i^{\check{m}-1}, o^{\check{m}-1}, j^2, p^2, \dots, j^{\underline{l}}, p^{\underline{l}})$$

(if $o^{\check{m}-1} = p^1$) is a CIC of μ for j^* at \hat{A} . However, if $m^* < \check{m}$, then

$$(i^{\check{m}}, o^{\check{m}}, \dots, i^{m(\bar{l})}, o^{m(\bar{l})}, j^{\bar{l}}, p^{\bar{l}}, \dots, j^{\underline{l}}, p^{\underline{l}}).$$

is a CIC of μ for \hat{j}^* at \hat{A} .

Next, consider the case where $p^{\underline{l}} \in A_{j^*}$. We have that either $p^{\underline{l}} = o^1$ or $p^{\underline{l}} \notin O(C)$. In both cases, $j^{\underline{l}+1} = j^*$. Since \hat{C} is a CIC for $\hat{j}^* > j^*$, we obtain $p^{\underline{l}+1} \in A_{j^*}$ as well.

We argue first that $m^* < \underline{m}$. If not, then

$$(j^1, p^1, \dots, j^{\underline{l}}, p^{\underline{l}}, i^{m(\underline{l})}, o^{m(\underline{l})}, \dots, i^{m(\bar{l})}, o^{m(\bar{l})}, j^{\bar{l}}, p^{\bar{l}}, \dots, j^{\underline{l}}, p^{\underline{l}})$$

is a CIC of μ for \hat{j}^* given that $m(\underline{l}) \leq m(\bar{l})$, $m(\bar{l}) < \bar{m}$, Claim 8, and $\{p^{\underline{l}}, p^{\underline{l}+1}\} \subseteq A_{j^*}$ jointly imply that there is no $m' \in \{m(\underline{l}), \dots, m(\bar{l})\}$ such that $i^{m'} < \hat{j}^*$ and such that C affects $i^{m'}$.

²¹Since $m(\underline{l}) < \underline{m}$, by definition of \underline{m} , either $m(\underline{l}) = 1$ or $i^{m(\underline{l})} \notin I(\hat{C})$. If $m(\underline{l}) \neq 1$ and $i^{m(\underline{l})} \notin I(\hat{C})$, then $p^{\underline{l}} = o^{m(\underline{l})-1}$. However, this implies $i^{m(\underline{l})-1} \in I(\hat{C})$. Since $m(\underline{l}) \neq 1$, this contradicts the definition of \underline{m} . Thus, $m(\underline{l}) = 1$ so either $p^{\underline{l}} = o^{m(\underline{l})-1} = o^M \in \mu(j^*)$ or $j^{\underline{l}^*} = j^*$ and $\underline{l} = \underline{l}^* - 1$ so that $p^{\underline{l}} \in (\mu + C)(j^*)$, which implies that $p^{\underline{l}} \in \mu(j^*)$ or $p^{\underline{l}} = o^1$.

Now let \tilde{l} be the first integer in $(\underline{l} + 1, \dots, \bar{l})$ such that either $j^{\tilde{l}} \in I(C) \setminus \{j^*\}$ or $p^{\tilde{l}} \in O(C)$. If $j^{\tilde{l}} \in I(C) \setminus \{j^*\}$, let \tilde{m} be such that $i^{\tilde{m}} = j^{\tilde{l}}$. Since $m^* < \underline{m}$,

$$(j^{l+1}, p^{l+1}, \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1}, i^{\tilde{m}}, o^{\tilde{m}}, \dots, i^M, o^M).$$

is a CIC of μ for j^* at \hat{A} . If $j^{\tilde{l}} \notin I(C) \setminus \{j^*\}$, we obtain a similar contradiction by letting \tilde{m} be such that $o^{\tilde{m}-1} = p^{\tilde{l}}$ and considering

$$(j^{l+1}, p^{l+1}, \dots, j^{\tilde{l}}, p^{\tilde{l}}, i^{\tilde{m}}, o^{\tilde{m}}, \dots, i^M, o^M).$$

Since we have now exhausted all the possibilities for the case of $m(\underline{l}) \leq m(\bar{l})$, we now consider the case of $m(\bar{l}) < m(\underline{l})$ for the remainder of the proof in Case 3. In this case, $\underline{m} \leq m(\underline{l})$.²²

If $\hat{j}^* \notin \{i^2, \dots, i^{m(\bar{l})}, i^{m(\underline{l})}, \dots, i^M\}$, then

$$(i^1, o^1, \dots, i^{m(\bar{l})}, o^{m(\bar{l})}, j^{\bar{l}}, p^{\bar{l}}, \dots, [p^L, j^1] \dots, j^L, p^L, i^{m(\underline{l})}, o^{m(\underline{l})}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at A that satisfies **(P1)** - **(P3)**.

If there exists a $\tilde{m} < m(\bar{l})$ such that $i^{\tilde{m}} = \hat{j}^*$ and $\{o^{\tilde{m}-1}, o^{\tilde{m}}\} \subseteq \Omega_{\hat{j}^*} \setminus A_{\hat{j}^*}$, then

$$(i^1, o^1, \dots, i^{\tilde{m}-1}, o^{\tilde{m}-1}, j^1, p^1, \dots, j^L, p^L, i^{m(\underline{l})}, o^{m(\underline{l})}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at A that satisfies **(P1)** - **(P3)**.

If there exists a $\tilde{m} \geq m(\underline{l})$ such that $i^{\tilde{m}} = \hat{j}^*$ and $\{o^{\tilde{m}-1}, o^{\tilde{m}}\} \subseteq \Omega_{\hat{j}^*} \setminus A_{\hat{j}^*}$, then

$$(j^1, p^1, \dots, j^L, p^L, i^{m(\underline{l})}, o^{m(\underline{l})}, \dots, i^{\tilde{m}-1}, o^{\tilde{m}-1})$$

is a CIC of μ for \hat{j}^* at \hat{A} since $m^* \notin \{\underline{m}, \dots, \tilde{m} - 1\}$.

If there exists a $\tilde{m} < m(\bar{l})$ such that $i^{\tilde{m}} = \hat{j}^*$ and $\{o^{\tilde{m}-1}, o^{\tilde{m}}\} \subseteq A_{\hat{j}^*}$, then

$$(i^{\tilde{m}}, o^{\tilde{m}}, \dots, i^{m(\bar{l})}, o^{m(\bar{l})}, j^{\bar{l}}, p^{\bar{l}}, \dots, j^L, p^L)$$

is a CIC of μ for \hat{j}^* at \hat{A} since $m^* \notin \{\underline{m}, \dots, \tilde{m} - 1\}$.

²²If $m(\bar{l}) < m(\underline{l}) < \underline{m}$, then $i^{m(\underline{l})} \notin I(\hat{C})$ and $m(\underline{l}) \neq 1$. So $p^L = o^{m(\underline{l})-1}$. This implies that $i^{m(\underline{l})-1} \in I(\hat{C})$. Since $m(\underline{l}) \neq 1$, this contradicts the definition of \underline{m} .

Finally, if there exists a $\tilde{m} \geq m(\underline{l})$ such that $i^{\tilde{m}} = \hat{j}^*$ and $\{o^{\tilde{m}-1}, o^{\tilde{m}}\} \subseteq A_{\hat{j}^*}$, then

$$(i^1, o^1, \dots, i^{m(\bar{l})}, o^{m(\bar{l})}, j^{\bar{l}}, p^{\bar{l}}, \dots, j^L, p^L, i^{\tilde{m}}, o^{\tilde{m}}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at A that satisfies (P1) - (P3).

Case 4: $\underline{m} < \bar{m}$, C does not affect \hat{j}^* , $p^L \notin O(C)$, and $j^{\bar{l}} = j^*$. By definition of $m(\bar{l})$ and the premise of this case, $m(\bar{l}) = M$. Hence, $m(\bar{l}) \geq m(\underline{l})$.

Assume first that $\{p^{\bar{l}-1}, p^{\bar{l}}\} \subseteq \Omega_{j^*} \setminus A_{j^*}$.

First, we claim that $\underline{l} < \bar{l} - 1$: otherwise, $\underline{l} = \bar{l} - 1$ and, given that $p^{\bar{l}-1} \notin O(C)$ (as $p^{\bar{l}-1} \neq o^1$) we have $\{p^1, \dots, p^{\bar{l}-1}\} \cap O(C) = \emptyset$; given that $p^L \notin O(C)$, Claim 6 and the definition of \bar{l} imply $\{p^{\bar{l}}, \dots, p^L\} \cap O(C) = \emptyset$, which contradicts Claim 4.

Second, we argue that $m(\underline{l}) \geq \underline{m}$: if $p^{\underline{l}} \notin O(C)$, we have $j^{\underline{l}+1} = i^{m(\underline{l})}$ and $\underline{l} < \bar{l} - 1$ implies $j^{\underline{l}+1} \neq j^*$ so $i^{m(\underline{l})-1} \in I(\hat{C}) \setminus \{j^*\}$; if $p^{\underline{l}} \in O(C)$, we have $j^{\underline{l}+1} = i^{m(\underline{l})-1}$ and $\underline{l} < \bar{l} - 1$ again implies $j^{\underline{l}+1} \neq j^*$ so $i^{m(\underline{l})-1} \in I(\hat{C}) \setminus \{j^*\}$.

Third, we show that $m^* < m(\underline{l})$: If not, Claim 7 and $m(\underline{l}) \geq \underline{m}$ imply $m^* \geq \bar{m}$. Let \tilde{l} be the last integer in $(1, \dots, \bar{l} - 1)$ such that $j^{\tilde{l}} \in I(C) \setminus \{j^*\}$ and let \tilde{m} be such that $i^{\tilde{m}} = j^{\tilde{l}}$. Since $p^{\tilde{l}-1} \notin O(C)$ in the case we consider here, we have $\{p^{\tilde{l}}, \dots, p^{\bar{l}-1}\} \cap O(C) = \emptyset$. But then $m^* \geq \bar{m} > \tilde{m} - 1$ implies that

$$(i^1, o^1, \dots, i^{\tilde{m}-1}, o^{\tilde{m}-1}, j^{\tilde{l}}, p^{\tilde{l}}, \dots, j^{\bar{l}-1}, p^{\bar{l}-1}).$$

is a CIC of μ for j^* at \hat{A} .

Now assume first that $\hat{j}^* \notin \{i^{m(\underline{l})}, \dots, i^M\}$. If $o^{m(\bar{l})} \neq p^{\bar{l}}$, then Claim 8 and $m^* < m(\underline{l})$ imply that

$$(j^1, p^1, \dots, j^{\underline{l}}, p^{\underline{l}}, i^{m(\underline{l})}, o^{m(\underline{l})}, \dots, i^{m(\bar{l})}, o^{m(\bar{l})}, j^{\bar{l}}, p^{\bar{l}}, \dots, j^L, p^L)$$

is a CIC of μ for \hat{j}^* at \hat{A} . If $o^{m(\bar{l})} = p^{\bar{l}}$, we obtain a contradiction to Claim 6 since $p^{\bar{l}} \in O(C)$ implies $\bar{l} = L$ but we have $p^L \notin O(C)$ in the case we consider here.

Next, assume that there exists a $\tilde{m} \in \{m(\underline{l}), \dots, M\}$ such that $i^{\tilde{m}} = \hat{j}^*$.

If $\{o^{\tilde{m}-1}, o^{\tilde{m}}\} \subseteq \Omega_{\hat{j}^*} \setminus A_{\hat{j}^*}$, then

$$(j^1, p^1, \dots, j^{\underline{l}}, p^{\underline{l}}, i^{m(\underline{l})}, o^{m(\underline{l})}, \dots, i^{\tilde{m}-1}, o^{\tilde{m}-1})$$

is a CIC of μ for \hat{j}^* at \hat{A} . If $\{o^{\tilde{m}-1}, o^{\tilde{m}}\} \subseteq A_{\hat{j}^*}$, then

$$(i^{\tilde{m}}, o^{\tilde{m}}, \dots, i^{m(\bar{l})}, o^{m(\bar{l})}, j^{\bar{l}}, p^{\bar{l}}, \dots, j^L, p^L)$$

is a CIC of μ for \hat{j}^* at \hat{A} .

We have now exhausted all the possibilities for the case of $\{o^{\tilde{m}-1}, o^{\tilde{m}}\} \subseteq \Omega_{\hat{j}^*} \setminus A_{\hat{j}^*}$. For the remainder of the proof in Case 4, we assume that $\{p^{\bar{l}-1}, p^{\bar{l}}\} \subseteq A_{\hat{j}^*}$.

If $\hat{j}^* \notin \{i^{m(\bar{l})}, \dots, i^M\}$, then

$$(j^{\bar{l}}, p^{\bar{l}}, \dots, [p^L, j^1] \dots, j^L, p^L, i^{m(\bar{l})}, o^{m(\bar{l})}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at A that satisfies (P1) - (P3).

Now assume that there exists $\tilde{m} \in \{m(\bar{l}), \dots, M\}$ such that $i^{\tilde{m}} = \hat{j}^*$.

If $\{o^{\tilde{m}-1}, o^{\tilde{m}}\} \subseteq \Omega_{\hat{j}^*} \setminus A_{\hat{j}^*}$, then, given that $\{p^{\bar{l}-1}, p^{\bar{l}}\} \subseteq A_{\hat{j}^*}$, the second part of Claim 8 implies that

$$(j^1, p^1, \dots, j^L, p^L, i^{m(\bar{l})}, o^{m(\bar{l})}, \dots, i^{\tilde{m}-1}, o^{\tilde{m}-1})$$

is a CIC of μ for \hat{j}^* at \hat{A} unless $m^* \in \{m(\bar{l}), \dots, \underline{m} - 1\}$. However, if $m^* < \underline{m}$, then

$$(j^{\bar{l}}, p^{\bar{l}}, \dots, j^L, p^L, i^{\tilde{m}}, o^{\tilde{m}}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at \hat{A} if $p^L \neq o^{\tilde{m}}$ and

$$(j^{\bar{l}}, p^{\bar{l}}, \dots, j^L, p^L, i^{\tilde{m}+1}, o^{\tilde{m}+1}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at \hat{A} if $p^L = o^{\tilde{m}}$.

Finally, if $\{o^{\tilde{m}-1}, o^{\tilde{m}}\} \subseteq A_{\hat{j}^*}$, then

$$(j^{\bar{l}}, p^{\bar{l}}, \dots, j^L, p^L, i^{\tilde{m}}, o^{\tilde{m}}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at A that satisfies (P1) - (P3).

Case 5: $\underline{m} < \bar{m}$, C does not affect \hat{j}^* , and $p^L \in O(C)$. Since $p^L \in O(C)$, $\hat{j}^* \in I(C)$.

Let \tilde{m} be such that $i^{\tilde{m}} = \hat{j}^*$. Since $p^L \in O(C)$, we have $p^L = o^{\tilde{m}}$ and, given that C does not affect \hat{j}^* , $o^{\tilde{m}-1} \in \Omega_{\hat{j}^*} \setminus A_{\hat{j}^*}$.

If $\tilde{m} < m(\bar{l})$, then

$$(i^1, o^1, \dots, i^{\tilde{m}-1}, o^{\tilde{m}-1}, j^1, p^1, \dots, j^L, p^L, i^{m(\bar{l})}, o^{m(\bar{l})}, \dots, i^M, o^M).$$

is a CIC of μ for j^* at A that satisfies (P1) - (P3).

Next, assume that $\check{m} = m(\underline{l})$. The definitions of \underline{l} and $m(\underline{l})$ imply that either $p^{\underline{l}} \in \mu(\hat{j}^*) \setminus (\mu + C)(\hat{j}^*)$ or that $\underline{l} = L, j^L \notin I(C)$, and $p^L \notin O(C)$. Since $p^L \in O(C)$ in the case we consider here, $p^{\underline{l}} = o^{\check{m}-1}$. However, given that $o \notin O(\hat{C})$ by Claim 7 and, by the premise of Case 5, $o^{\check{m}-1} = p^{\underline{l}} \in \Omega_{j^*} \setminus A_{j^*}$, we obtain that

$$(j^1, p^1, \dots, j^{\underline{l}}, p^{\underline{l}})$$

is a CIC of μ for \hat{j}^* at \hat{A} .

For the remainder of the proof in Case 5, we assume that $\check{m} > m(\underline{l})$.

We argue first that $m(\underline{l}) < \underline{m}$: Otherwise, Claim 7 implies $m^* \notin \{m(\underline{l}), \dots, \check{m} - 1\}$ and Claim 8 implies that

$$(j^1, p^1, \dots, j^{\underline{l}}, p^{\underline{l}}, i^{m(\underline{l})}, o^{m(\underline{l})}, \dots, i^{\check{m}-1}, o^{\check{m}-1})$$

is a CIC of μ for \hat{j}^* at \hat{A} .

As for Case 3, $m(\underline{l}) < \underline{m}$ implies that either $p^{\underline{l}} \in \mu(j^*)$ or $p^{\underline{l}} = o^1$.

If $p^{\underline{l}} \in \Omega_{j^*} \setminus A_{j^*}$, then

$$(j^1, p^1, \dots, j^{\underline{l}}, p^{\underline{l}}, i^1, o^1, \dots, i^{\check{m}-1}, o^{\check{m}-1})$$

is a CIC of μ for j^* at A that satisfies (P1) - (P3).

If $p^{\underline{l}} \in A_{j^*}$, then $j^{\underline{l}+1} = j^*$.

We argue first that we must have $m^* < \underline{m}$: Otherwise, either

$$(j^1, p^1, \dots, j^{\underline{l}}, p^{\underline{l}}, i^1, o^1, \dots, i^{\check{m}-1}, o^{\check{m}-1})$$

(if $p^{\underline{l}} \neq o^1$) or

$$(j^1, p^1, \dots, j^{\underline{l}}, p^{\underline{l}}, i^2, o^2, \dots, i^{\check{m}-1}, o^{\check{m}-1})$$

(if $p^{\underline{l}} = o^1$) would be a CIC of μ for \hat{j}^* at \hat{A} .

Next, since \hat{C} is a CIC for $\hat{j}^* > j^*$, we have $p^{\underline{l}+1} \in A_{j^*}$ as well. Let \tilde{l} be the first integer in $(\underline{l} + 1, \dots, \bar{l})$ such that either $j^{\tilde{l}} \in I(C) \setminus \{j^*\}$ or $p^{\tilde{l}} \in O(C)$. If $j^{\tilde{l}} \in I(C) \setminus \{j^*\}$, let \tilde{m} be such that $i^{\tilde{m}} = j^{\tilde{l}}$. Since $m^* < \underline{m}$,

$$(j^{\underline{l}+1}, p^{\underline{l}+1}, \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1}, i^{\tilde{m}}, o^{\tilde{m}}, \dots, i^M, o^M).$$

is a CIC of μ for j^* at \hat{A} . If $j^{\bar{l}} \notin I(C) \setminus \{j^*\}$, we obtain a similar contradiction by letting \tilde{m} be such that $o^{\tilde{m}-1} = p^{\bar{l}}$ and considering

$$(j^{l+1}, p^{l+1}, \dots, j^{\bar{l}}, p^{\bar{l}}, i^{\tilde{m}}, o^{\tilde{m}}, \dots, i^M, o^M).$$

□

Proof of Theorem 3. Let A be a profile of desirable sets and, for some $i \in I$, let \hat{A}_i be an alternative report for i .

We argue first that it is without loss of generality to assume that $\hat{A}_i \setminus A_i = \emptyset$. To see this, assume that $\hat{A}_i \setminus A_i \neq \emptyset$ and let $o \in \hat{A}_i \setminus A_i$. First suppose that $o \notin \Omega_i \cup A_i$. If $K^i(\hat{A}_i \setminus \{o\}) = K^i(\hat{A}_i)$, then it is immediate that we can consider the manipulation $\hat{A}_i \setminus \{o\}$ rather than \hat{A}_i . If $K^i(\hat{A}_i \setminus \{o\}) \neq K^i(\hat{A}_i)$, then Lemma 1 implies that $o \in \mu(i)$ for all $\mu \in \mathcal{M}^i(\hat{A})$. Since $o \in \hat{A}_i \setminus (\Omega_i \cup A_i)$, each matching in $\mathcal{M}^i(\hat{A})$ is unacceptable to i (w.r.t. her true preferences). Now suppose that $o \in \Omega_i$. Then by Lemma 2, for each $\mu \in \mathcal{M}^i(\hat{A})$ and each $\mu' \in \mathcal{M}^i(A)$, $|\mu'(i) \cap A_i| = K^i(A) \geq |\mu(i) \cap A_i|$. Hence, reporting $\hat{A}'_i \equiv \hat{A}_i \setminus \{o\}$ is at least as good for i as \hat{A}_i when her true desirable set is A_i . Applying these arguments $|\hat{A}_i \setminus A_i|$ times, we end up with a manipulation \hat{A}'_i that does at least as well as \hat{A}_i and that satisfies $\hat{A}'_i \subseteq A_i$.

We now consider \hat{A}_i such that $\hat{A}_i \subsetneq A_i$. Let $\{o^1, \dots, o^T\} = A_i \setminus \hat{A}_i$ for some $T \geq 1$. By Lemma 3 (taking $\hat{A}_i \cup \{o^1\}$ to be the true and \hat{A}_i to be the false set of desirable objects), we obtain $K^i(\hat{A}_i \cup \{o^1\}) \geq K^i(\hat{A}_i)$. Proceeding inductively, T applications of Lemma 3 yield that $K^i(\hat{A}_i \cup \{o^1, \dots, o^T\}) \geq K^i(\hat{A}_i)$. Hence, by definition of o^1, \dots, o^M we obtain that $K^i(A) \geq K^i(\hat{A}_i)$. This completes the proof.

□

D Claims in Proof of Lemma 3

Proof of Claim 1. Since $\mu \in \mathcal{M}^{i-1}(\hat{A})$ and $j^* < i$, we obtain $\mu(j^*) \cap \hat{A}_{j^*} = K^{j^*}(\hat{A})$. Since $K^{j^*}(A) > K^{j^*}(\hat{A})$, we obtain $\mu \notin \mathcal{M}^{j^*}(A)$.

On the other hand, $j^* < i$ also implies that $\mu \in \mathcal{M}^{j^*-1}(\hat{A})$ given that $\mathcal{M}^{i-1}(\hat{A}) \subseteq \mathcal{M}^{j^*-1}(\hat{A})$. Since $\hat{A}_j \subseteq A_j$ for all $j \in I$ and $K^j(\hat{A}) = K^j(A)$ if $j < j^*$, the definition of j^* implies that $\mathcal{M}^{j^*-1}(\hat{A}) \subseteq \mathcal{M}^{j^*-1}(A)$ and thus $\mu \in \mathcal{M}^{j^*-1}(A)$. □

Proof of Claim 2. If not, then $\mu + C \in \mathcal{M}^{j^*-1}(\hat{A})$ since $\mu \in \mathcal{M}^{j^*-1}(\hat{A})$, $o \notin (\mu + C)(i)$, $\hat{A}_i = A_i \setminus \{o\}$, and $\hat{A}_j = A_j$ for all $j \neq i$. But $\mu + C \in \mathcal{M}^{j^*-1}(\hat{A})$ contradicts $\mu \in \mathcal{M}^{i-1}(\hat{A}) \subseteq \mathcal{M}^{j^*}(\hat{A})$ since $|(\mu + C)(j^*) \cap \hat{A}_{j^*}| > |\mu(j^*) \cap \hat{A}_{j^*}|$. □

Proof of Claim 3. Note that the first part of Theorem 2 implies that $\hat{j}^* \geq j^*$ since $\mu + C \in \mathcal{M}^{j^*-1}(A)$ given that $\mu \in \mathcal{M}^{j^*-1}(A)$ and that C is a CIC of μ for j^* at A . Given that $\hat{j}^* \geq j^*$ and $\mu \in \mathcal{M}^{j^*-1}(A)$, our assumption that \hat{j}^* is the highest priority agent for whom there exists a CIC of $\mu + C$ at A and the second part of Theorem 2 imply $\mu + C \in \mathcal{M}^{\hat{j}^*-1}(A)$. Furthermore, we also have $(\mu + C) + \hat{C} \in \mathcal{M}^{\hat{j}^*-1}(A)$ since $\mu + C \in \mathcal{M}^{\hat{j}^*-1}(A)$ and since \hat{C} is a CIC of $\mu + C$ for \hat{j}^* at A . \square

Proof of Claim 4. If $O(C) \cap O(\hat{C}) = \emptyset$, then by Claim 2, $o \notin O(\hat{C})$. Furthermore, $O(C) \cap O(\hat{C}) = \emptyset$ also implies that for all $j \in I$ and all $o' \in O(\hat{C})$, $o' \in (\mu + C)(j)$ if and only if $o' \in \mu(j)$. Combining the last two observations, we find that \hat{C} is a CIC of μ for \hat{j}^* at \hat{A} . Since $\hat{j}^* < i$, this contradicts the assumption that $\mu \in \mathcal{M}^{i-1}(\hat{A}) \subseteq \mathcal{M}^{\hat{j}^*}(\hat{A})$. \square

Proof of Claim 5. Assume to the contrary that there is an agent $\underline{j} < j^*$ for whom the statement of the claim is violated. Without loss of generality suppose that there is no agent with higher priority than \underline{j} for whom Claim 5 does not hold, that is, for all pairs m', l' such that $i^{m'} = j^{l'}$ and $i^{m'} < \underline{j}$, either

$$\{o^{m'-1}, o^{m'}, p^{l'-1}, p^{l'}\} \subseteq A_{i^{m'}} \text{ or } \{o^{m'-1}, o^{m'}, p^{l'-1}, p^{l'}\} \subseteq \Omega_{i^{m'}} \setminus A_{i^{m'}}.$$

Since C and \hat{C} are CICs for agents with lower priority than \underline{j} , there are only two cases to consider. In either case, we construct a CIC of μ for \underline{j} at A . Since $\underline{j} < j^*$, by Theorem 2, the existence of such a CIC contradicts the implication of Claim 2 that $\mu \in \mathcal{M}^{j^*-1}(A)$.

Case 1: $\{o^{m-1}, o^m\} \subseteq \Omega_{\underline{j}} \setminus A_{\underline{j}}$ and $\{p^{l-1}, p^l\} \subseteq A_{\underline{j}}$. Let \tilde{l} be the first integer in $(l, \dots, [L, 1], \dots, l-1)$ such that either $j^{\tilde{l}} \in I(C) \setminus \{\underline{j}\}$ or $p^{\tilde{l}} \in O(C)$. Note that the existence of an \tilde{l} with the desired properties follows since $O(C) \cap O(\hat{C}) \neq \emptyset$ by Claim 4. If $j^{\tilde{l}} \in I(C) \setminus \{\underline{j}\}$, let \tilde{m} be such that $i^{\tilde{m}} = j^{\tilde{l}}$ and note that our assumption about \underline{j} implies that

$$(j^l, p^l, \dots, [p^L, j^1], \dots, j^{\tilde{l}}, p^{\tilde{l}}, i^{\tilde{m}}, o^{\tilde{m}}, \dots, [o^M, i^1], \dots, i^{m-1}, o^{m-1})$$

is a CIC of μ for \underline{j} at A .

If $j^{\tilde{l}} \notin I(C) \setminus \{\underline{j}\}$ and $p^{\tilde{l}} \notin \mu(\underline{j})$, let \tilde{m} be such that $o^{\tilde{m}} = p^{\tilde{l}}$ and note that our assumption about \underline{j} implies that

$$(j^l, p^l, \dots, [p^L, j^1], \dots, j^{\tilde{l}}, p^{\tilde{l}}, i^{\tilde{m}+1}, o^{\tilde{m}+1}, \dots, [o^M, i^1], \dots, i^{m-1}, o^{m-1})$$

is a CIC of μ for \underline{j} at A .

Finally, if $j^{\bar{l}} \notin I(C) \setminus \{\underline{j}\}$ and $p^{\bar{l}} \in \mu(\underline{j})$, then

$$(j^{\bar{l}}, p^{\bar{l}}, \dots [p^L, j^1] \dots, j^{\bar{l}}, p^{\bar{l}})$$

is a CIC of μ for \underline{j} at A since $p^{\bar{l}} = o^{m-1} \in \Omega_{\underline{j}} \setminus A_{\underline{j}}$.

Case 2: $\{o^{m-1}, o^m\} \subseteq A_{\underline{j}}$ and $\{p^{l-1}, p^l\} \subseteq \Omega_{\underline{j}} \setminus A_{\underline{j}}$. Let \tilde{l} be the last integer in $(l, \dots [L, 1] \dots, l-1)$ such that either $j^{\tilde{l}} \in I(C) \setminus \{\underline{j}\}$ or $p^{\tilde{l}} \in O(C)$. Claim 4 ensures that such \tilde{l} exists.

We now establish that $p^{\tilde{l}} \notin O(C)$. If there is \tilde{m} such that $p^{\tilde{l}} = o^{\tilde{m}}$, then since $p^{\tilde{l}} \in (\mu + C)(j^{\tilde{l}+1})$, we have $j^{\tilde{l}+1} = i^{\tilde{m}} \in I(C)$. Since $j^{\tilde{l}+1} \in I(C)$ is a contradiction to the definition of \tilde{l} if $\tilde{l} \neq l-1$, we have $\tilde{l} = l-1$. Given that $p^{l-1} \in (\mu + C)(\underline{j})$ and C is a CIC of μ , $p^{l-1} \in O(C)$ is possible only if $p^{l-1} = o^m$. However, since $o^m \in A_{\underline{j}}$ and $p^{l-1} \in \Omega_{\underline{j}} \setminus A_{\underline{j}}$, we have $o^m \neq p^{l-1}$ and hence, by our previous arguments, $p^{l-1} \notin O(C)$. Since $p^{\tilde{l}} \notin O(C)$, we have $j^{\tilde{l}} \in I(C)$ so that there is an \hat{m} be such that $i^{\hat{m}+1} = j^{\tilde{l}}$.

By an argument analogous to that in Case 1, the following is a CIC of μ for \underline{j} at A :

$$\tilde{C} \equiv (i^{\hat{m}}, o^{\hat{m}}, \dots [o^M, i^1] \dots, i^{\hat{m}}, o^{\hat{m}}, j^{\tilde{l}}, p^{\tilde{l}}, \dots [p^L, j^1] \dots, j^{l-1}, p^{l-1})$$

□

Proof of Claim 6. Suppose that $p^{\bar{l}} \in O(C)$. Since \hat{C} is a CIC of $\mu + C$, we have $j^{\bar{l}+1} \in I(C)$. However, if $\bar{l} < L$, then $\bar{l} + 1 \neq 1$, so $j^{\bar{l}+1} \neq j^1 = \hat{j}^*$. This contradicts the definition of \bar{l} . □

Proof of Claim 7. We argue first that if $\underline{m} \leq m^*$, then $o \notin O(\hat{C})$ and $\bar{m} \leq m^*$.

Assume to the contrary that $\underline{m} \leq m^*$ and either $o \in O(\hat{C})$ or $m^* < \bar{m}$. Let $\tilde{m}_1, \dots, \tilde{m}_T$ be such that

1. $i^{\tilde{m}_t} \in I(\hat{C}) \setminus \{j^*\}$ for all $t \in \{1, \dots, T\}$
2. $\tilde{m}_t < \tilde{m}_{t+1}$ for all $t \leq T-1$ and $\tilde{m}_T \leq m^*$
3. For all $m' \in \{2, \dots, m^*\} \setminus \{\tilde{m}_1, \dots, \tilde{m}_T\}$, $i^{m'} \notin I(\hat{C})$

Let $\tilde{l}_1, \dots, \tilde{l}_T$ be such that $j^{\tilde{l}_t} = i^{\tilde{m}_t}$ for all $t \in \{1, \dots, T\}$. Note that since $\underline{m} < m^*$ and either $o^{m^*} = o \in O(\hat{C})$ or $m^* < \bar{m}$, there exists $t^* \leq T$ such that

$$\{p^{\tilde{l}_{t^*}}, j^{\tilde{l}_{t^*}+1}, p^{\tilde{l}_{t^*}+1}, \dots [p^L, j^1] \dots, j^{\tilde{l}_{t^*}+1-1}, p^{\tilde{l}_{t^*}+1-1}\} \cap \{o^{m^*}, i^{m^*+1}, o^{m^*+1}, \dots, i^M, o^M\} \neq \emptyset, \quad (4)$$

where $\tilde{l}_{t^*+1} = \tilde{l}_1$ if $t^* = T$. For the remainder of the proof of Claim 7, fix some t^* for which Equation (4) holds and let \hat{l} be the first integer in $(\tilde{l}_{t^*}, \dots [L, 1] \dots, \tilde{l}_{t^*+1} - 1)$ such that either $p^{\hat{l}} \in \{o^{m^*}, \dots, o^M\}$ or $j^{\hat{l}} \in \{i^{m^*+1}, \dots, i^M\}$.

We distinguish two cases:

Case 1: $\hat{l} \neq \tilde{l}_{t^*}$ and $j^* \in \{j^{\tilde{l}_{t^*}+1}, \dots [j^L, j^1] \dots, j^{\hat{l}-1}\}$.

Assume first that $j^* = \hat{j}^* = j^1$. If $j^{\hat{l}} \in \{i^{m^*+1}, \dots, i^M\}$, let \hat{m} be such that $i^{\hat{m}} = j^{\hat{l}}$ and consider the sequence

$$\tilde{C} \equiv (j^1, p^1, \dots, j^{\hat{l}-1}, p^{\hat{l}-1}, i^{\hat{m}}, o^{\hat{m}}, \dots, i^M, o^M).$$

Note that $\hat{l} > 1$ since $j^{\hat{l}} \in \{i^{m^*+1}, \dots, i^M\}$ and $j^1 = i^1 = j^*$ in the case we consider here. By the definitions of \tilde{l}_{t^*} and \tilde{l}_{t^*+1} as well as the fact that $\hat{l} \leq \tilde{l}_{t^*+1} - 1$, we obtain $\{p^1, \dots, p^{\hat{l}-1}\} \cap O(C) = \emptyset$. Since $p^1 \in A_{j^*}$ and $o^M \in \Omega_{j^*} \setminus A_{j^*}$, Claim 5 implies that \tilde{C} is a CIC of μ for j^* at \hat{A} , which is a contradiction to $\mu \in \mathcal{M}^{i^{-1}}(\hat{A}) \subseteq \mathcal{M}^{j^*}(\hat{A})$. If $p^{\hat{l}} \in \{o^{m^*}, \dots, o^M\}$ and $j^{\hat{l}} \notin \{i^{m^*+1}, \dots, i^M\}$, let \hat{m} be such that $o^{\hat{m}} = p^{\hat{l}}$ and consider

$$(j^1, p^1, \dots, j^{\hat{l}}, p^{\hat{l}}, i^{\hat{m}+1}, o^{\hat{m}+1}, \dots, i^M, o^M)$$

to obtain a contradiction in a similar manner as before.²³

Henceforth, we assume that $j^* \neq \hat{j}^*$. Let $\tilde{l} \geq 2$ be such that $j^{\tilde{l}} = j^*$ and note that Claim 3 implies $j^{\tilde{l}} < \hat{j}^*$.

Assume first that $p^{\tilde{l}} \in A_{j^*}$. If $j^{\hat{l}} \in \{i^{m^*+1}, \dots, i^M\}$, then $\hat{l} \neq \tilde{l}$ since $j^{\tilde{l}} = j^* = i^1$. Analogs of the arguments above imply that

$$(j^{\tilde{l}}, p^{\tilde{l}}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1}, i^{\hat{m}}, o^{\hat{m}}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at \hat{A} , which contradicts $\mu \in \mathcal{M}^{i^{-1}}(\hat{A}) \subseteq \mathcal{M}^{j^*}(\hat{A})$. If $p^{\tilde{l}} \in \{o^{m^*}, \dots, o^M\}$ and $j^{\hat{l}} \notin \{i^{m^*+1}, \dots, i^M\}$, let \hat{m} be such that $o^{\hat{m}} = p^{\tilde{l}}$ and consider

$$\tilde{C}' \equiv (j^{\tilde{l}}, p^{\tilde{l}}, \dots [p^L, j^1] \dots, j^{\hat{l}}, p^{\hat{l}}, i^{\hat{m}+1}, o^{\hat{m}+1}, \dots, i^M, o^M).²⁴$$

Since i does not point to o in \tilde{C}' , analogs of our earlier arguments imply that \tilde{C}' is a CIC of μ at \hat{A} , which is impossible given our choice of μ .

²³The only difference to before is that the sequence may now contain o . However, given that $\hat{m} + 1 > m^*$, it cannot be the case that i points to o in the proposed cycle.

²⁴Note that $\{i^{\hat{m}+1}, o^{\hat{m}+1}, \dots, i^M, o^M\} = \emptyset$ if $\hat{m} = M$.

Next, consider the case where $p^{\tilde{l}} \in \Omega_{j^*} \setminus A_{j^*}$. Since $j^* < \hat{j}^*$ and since \hat{C} is a CIC of $\mu + C$ for \hat{j}^* , we have $p^{\tilde{l}-1} \in \Omega_{j^*} \setminus A_{j^*}$ as well. Consider the sequence

$$\tilde{C}'' \equiv (i^1, o^1, \dots, i^{\tilde{m}_{t^*}-1}, o^{\tilde{m}_{t^*}-1}, j^{\tilde{l}_{t^*}}, p^{\tilde{l}_{t^*}}, \dots [p^L, j^1] \dots j^{\tilde{l}-1}, p^{\tilde{l}-1}).$$

Note that $\tilde{m}_{t^*} \geq 2$ and $\tilde{l}_{t^*} \neq \tilde{l}$ since $i^{\tilde{m}_{t^*}} \in I(C) \setminus \{j^*\}$ and $j^* = j^{\tilde{l}}$. We now show that $\{p^{\tilde{l}_{t^*}}, \dots [p^L, p^1] \dots, p^{\tilde{l}-1}\} \cap O(C) = \emptyset$. If $p^l \in O(C)$ for some $l \in \{\tilde{l}_{t^*}, \dots [L, 1] \dots, \tilde{l}-2\}$, then, since \hat{C} is a CIC of $\mu + C$, we have $j^{l+1} \in I(C)$. Since $l \in \{l_{t^*}+1, \dots [L, 1] \dots, l_{t^*}+1-1\}$, the construction of the sequence $\{\tilde{l}_t\}_{t=1}^T$ implies that $j^{l+1} \in \{i^{m^*+1}, \dots, i^M\}$. However, since $\tilde{l} \in \{\tilde{l}_{t^*} + 1, \dots [L, 1] \dots, \tilde{l} - 1\}$ and $\{p^{\tilde{l}_{t^*}}, j^{\tilde{l}_{t^*}+1}, p^{\tilde{l}_{t^*}+1}, \dots [p^L, j^1] \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1}\} \cap \{o^{m^*}, i^{m^*+1}, o^{m^*+1}, \dots, i^M, o^M\} = \emptyset$, $j^{l+1} \in \{i^{m^*+1}, \dots, i^M\}$ is impossible. If $p^{\tilde{l}-1} \in O(C)$, then, since \hat{C} is a CIC of $\mu + C$, we have $p^{\tilde{l}-1} = o^1$. However, since $p^{\tilde{l}-1} \in \Omega_{j^*} \setminus A_{j^*}$ and $o^1 \in A_{j^*}$ we have $p^{\tilde{l}-1} \neq o^1$. Given that we have now established $\{p^{\tilde{l}_{t^*}+1}, \dots [p^L, p^1] \dots, p^{\tilde{l}-1}\} \cap O(C) = \emptyset$, similar arguments as before show that \tilde{C}'' satisfies all conditions of a CIC of μ for j^* at A . Furthermore, since $\tilde{m}_{t^*} - 1 < m^*$, we have that $o \notin O(\tilde{C}'')$ and \tilde{C}'' is also a CIC of μ for j^* at \hat{A} , which is a contradiction.

Case 2: $\hat{l} = \tilde{l}_{t^*}$ or $\hat{l} \neq \tilde{l}_{t^*}$ and $j^* \notin \{j^{\tilde{l}_{t^*}+1}, \dots [j^L, j^1] \dots, j^{\hat{l}-1}\}$.

Let $\tilde{m} \equiv \tilde{m}_{t^*}$, $\tilde{l} \equiv \tilde{l}_{t^*}$, and \hat{m} be such that $i^{\hat{m}} = j^{\hat{l}}$ if $j^{\hat{l}} \in I(C)$ and $o^{\hat{m}} = p^{\hat{l}}$ if $j^{\hat{l}} \notin I(C)$. Note that the properties that define $\tilde{m}, \tilde{l}, \hat{l}, \hat{m}$ imply that if $p^{\hat{l}} \notin \{o^{m^*}, \dots, o^M\}$, then $\hat{l} \neq \tilde{l}$ since $\tilde{m} = \tilde{m}_{t^*} \leq m^*$.

If $p^{\hat{l}} \in \{o^{m^*}, \dots, o^M\}$ and $j^{\hat{l}} \notin I(C)$, then we claim that

$$\tilde{C} \equiv (i^1, o^1, \dots, i^{\tilde{m}-1}, o^{\tilde{m}-1}, j^{\tilde{l}}, p^{\tilde{l}}, \dots [p^L, j^1] \dots, j^{\hat{l}}, p^{\hat{l}}, i^{\hat{m}+1}, o^{\hat{m}+1}, \dots i^M, o^M)$$

is a CIC of μ for j^* at A .²⁵ The arguments to show that \tilde{C} is a cycle of μ are analogous to the corresponding arguments in the proof of Claim 5. Next, recall that, by Claim 3, $j^* \leq \hat{j}^*$. Hence, Claim 5 implies that \tilde{C} does not affect any agent $k < j^* = \min\{j^*, \hat{j}^*\}$. Since $o^1 \in A_{j^*}$ as well as $o^M \in \Omega_{j^*} \setminus A_{j^*}$, and since $i^1 = j^*$ is the only instance of j^* in \tilde{C} , \tilde{C} increases the welfare of j^* . Note that the definitions of $\tilde{m}, \tilde{l}, \hat{l}, \hat{m}$ imply that $i \in I(\tilde{C})$ only if $i = j^{\tilde{l}}$. However, since \hat{C} is a CIC of $\mu + C$ and $o \in (\mu + C)(i) \setminus \mu(i)$, we have $p^{\tilde{l}} \neq o$. Hence, $o \notin (\mu + \tilde{C})(i)$ and therefore \tilde{C} is a CIC of μ for j^* at \hat{A} as well, contradicting the assumption that $\mu \in \mathcal{M}^{i-1}(\hat{A}) \subseteq \mathcal{M}^{j^*}(\hat{A})$.

²⁵Note that $\tilde{m} \geq 2$ and that $\{i^{\hat{m}+1}, o^{\hat{m}+1}, \dots i^M, o^M\} = \emptyset$ if $p^{\hat{l}} = o^M$. Note also that $o^M \in O(\tilde{C})$.

If $j^{\hat{l}} \in I(C)$, then analogous arguments establish that

$$(i^1, o^1, \dots, i^{\hat{m}-1}, o^{\hat{m}-1}, j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1}, i^{\hat{m}}, o^{\hat{m}}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at \hat{A} and we again obtain a contradiction.

We have thus shown that if $\underline{m} \leq m^*$, then $o \notin O(\hat{C})$ and $\bar{m} \leq m^*$.

Finally, we now show that $o \notin O(\hat{C})$. Assume to the contrary that there exists an l such that $p^l = o$. Since \hat{C} is a CIC of $\mu + C$ at A and $o^{m^*} = o$, we have that $j^{l+1} = i^{m^*} = i$. Since $i \in I(\hat{C})$, $\underline{m} \leq m^*$ so that our previous arguments imply $o \notin O(\hat{C})$, thus contradicting our hypothesis that $o \in O(\hat{C})$. \square

Proof of Claim 8. We proceed by contradiction. Without loss of generality, assume that \check{m} is such that $i^{\check{m}}$ is highest priority agent for whom one of the conditions in Claim 8 is violated. Whenever $i^{\check{m}} \in I(\hat{C})$, we let \check{l} be such that $j^{\check{l}} = i^{\check{m}}$. Furthermore, whenever $j^* \in I(\hat{C})$, we let l^* be such that $j^{l^*} = j^*$. Since we only work with the reassignments implied by C and \hat{C} , Claim 5 implies that the CICs we construct cannot affect any agent $j' < j^*$. We use this observation repeatedly throughout our proof of Claim 8.

We distinguish 12 cases to cover all the ways in which the conditions of Claim 8 may be violated for $i^{\check{m}}$. Below is a roadmap that explains which cases cover each possible violation.

- If Part (i) of the first statement is violated, then Case 1 or 2 applies.
- If Part (ii) of the first statement is violated, then Case 11 or 12 applies.
- If Part (i) of the second statement is violated, then Case 5 or 6 applies.
- If Part (ii) of the second statement is violated, then Case 3, 7, 11, or 12 applies.
- If Part (i) of the third statement is violated, then Case 8 or 9 applies.
- If Part (ii) of the third statement is violated, then Case 4, 10, 11, or 12 applies.
- If the fourth statement is violated, then Case 1, 3, or 4 applies.²⁶

Case 1: $\check{m} \in \{\underline{m} + 1, \dots, \bar{m} - 1\}$, $i^{\check{m}} \leq \hat{j}^*$, $o^{\check{m}-1} \in \Omega_{i^{\check{m}}} \setminus A_{i^{\check{m}}}$, and $o^{\check{m}} \in A_{i^{\check{m}}}$. By our assumption about \check{m} , there exist \hat{m}_1 , \hat{m}_2 , \hat{l}_1 , and \hat{l}_2 such that

$$1. \underline{m} \leq \hat{m}_1 < \check{m} < \hat{m}_2 \leq \bar{m}$$

²⁶Note that since \hat{C} is a CIC of $\mu + C$ for \hat{j}^* , $i^{\check{m}} = \hat{j}^*$ implies $\check{m} \in \{\underline{m}, \dots, \bar{m}\}$ and that \hat{C} increases the welfare of $i^{\check{m}}$.

2. $i^{\hat{m}_1} = j^{\hat{l}_1}$ and $i^{\hat{m}_2} = j^{\hat{l}_2}$
3. $\{j^{\hat{l}_2+1}, \dots [j^L, j^1] \dots, j^{\hat{l}_1-1}\} \cap I(C) \subseteq \{i^{\hat{m}}, j^*\}$

Assume first that $i^{\hat{m}} \notin \{j^{\hat{l}_2+1}, \dots [j^L, j^1] \dots, j^{\hat{l}_1-1}\}$ and that $o^1 \notin \{p^{\hat{l}_2}, \dots [p^L, p^1] \dots, p^{\hat{l}_1-1}\}$. If $p^{\hat{l}_1-1} \notin O(C)$, then

$$(i^{\hat{m}_1}, o^{\hat{m}_1}, \dots, i^{\hat{m}_2-1}, o^{\hat{m}_2-1}, j^{\hat{l}_2}, p^{\hat{l}_2}, \dots [p^L, j^1] \dots, j^{\hat{l}_1-1}, p^{\hat{l}_1-1})$$

is a CIC of μ for $i^{\hat{m}}$ at A . Since Claim 7 implies that $m^* \notin \{\hat{m}_1, \dots, \hat{m}_2 - 1\}$, the preceding sequence is a CIC of μ for $i^{\hat{m}}$ at \hat{A} , which contradicts $\mu \in \mathcal{M}^{i^{-1}}(\hat{A})$. If $p^{\hat{l}_1-1} \in O(C)$, then $p^{\hat{l}_1-1} = o^{\hat{m}_1}$ and so we obtain a similar contradiction using

$$(i^{\hat{m}_1+1}, o^{\hat{m}_1+1}, \dots, i^{\hat{m}_2-1}, o^{\hat{m}_2-1}, j^{\hat{l}_2}, p^{\hat{l}_2}, \dots [p^L, j^1] \dots, j^{\hat{l}_1-1}, p^{\hat{l}_1-1}).$$

Next, assume that $i^{\hat{m}} \notin \{j^{\hat{l}_2+1}, \dots [j^L, j^1] \dots, j^{\hat{l}_1-1}\}$ and that there exists $l^* \in \{\hat{l}_2 + 1, \dots [L, 1] \dots, \hat{l}_1\}$ such that $p^{l^*-1} = o^1$. Since $o^1 \in A_{j^*}$, we must have $j^* < \hat{j}^*$ and since \hat{C} is a CIC for \hat{j}^* , we obtain $p^{l^*} \in A_{j^*}$. If $m^* < \underline{m}$ and $p^{\hat{l}_1-1} \notin O(C)$, then

$$(j^{l^*}, p^{l^*}, \dots [p^L, j^1] \dots, j^{\hat{l}_1-1}, p^{\hat{l}_1-1}, i^{\hat{m}_1}, o^{\hat{m}_1}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at \hat{A} . If $m^* < \underline{m}$ and $p^{\hat{l}_1-1} = o^{\hat{m}_1}$, we obtain a similar contradiction by considering

$$(j^{l^*}, p^{l^*}, \dots [p^L, j^1] \dots, j^{\hat{l}_1-1}, p^{\hat{l}_1-1}, i^{\hat{m}_1+1}, o^{\hat{m}_1+1}, \dots, i^M, o^M).$$

If $m^* \geq \underline{m}$, then Claim 7 implies $m^* \geq \overline{m}$ and

$$(i^2, o^2, \dots, i^{\hat{m}_2-1}, o^{\hat{m}_2-1}, j^{\hat{l}_2}, p^{\hat{l}_2}, \dots [p^L, j^1] \dots, j^{l^*-1}, p^{l^*-1})$$

is a CIC of μ for $i^{\hat{m}}$ at \hat{A} .

Next, assume that there exists $\check{l} \in \{\hat{l}_2 + 1, \dots [L, 1] \dots, \hat{l}_1 - 1\}$ such that $j^{\check{l}} = i^{\hat{m}}$.

If $p^{\check{l}} \in A_{j^i}$ and $o^1 \notin \{p^{\check{l}}, \dots [p^L, p^1] \dots, p^{\hat{l}_1-1}\}$, then, given that $m^* \notin \{\underline{m}, \dots, \overline{m} - 1\}$,

$$(j^{\check{l}}, p^{\check{l}}, \dots [p^L, j^1] \dots, j^{\hat{l}_1-1}, p^{\hat{l}_1-1}, i^{\hat{m}_1}, o^{\hat{m}_1}, \dots, i^{\hat{m}-1}, o^{\hat{m}-1})$$

is a CIC of μ for $j^{\check{l}}$ at \hat{A} if $p^{\hat{l}_1-1} \notin O(C)$ and

$$(j^{\check{l}}, p^{\check{l}}, \dots [p^L, j^1] \dots, j^{\hat{l}_1-1}, p^{\hat{l}_1-1}, i^{\hat{m}_1+1}, o^{\hat{m}_1+1}, \dots, i^{\hat{m}-1}, o^{\hat{m}-1})$$

is a CIC of μ for $j^{\tilde{l}}$ at \hat{A} if $p^{\hat{l}_1-1} = o^{\hat{m}_1}$. Next, assume that $p^{\tilde{l}} \in A_{j^{\tilde{l}}}$ and there is $l^* \in \{\tilde{l} + 1, \dots [L, 1] \dots, \hat{l}_1\}$ such that $p^{l^*-1} = o^1$. If $m^* < \underline{m}$, we obtain a contradiction just as in the case where $i^{\tilde{m}} \notin \{j^{\hat{l}_2}, \dots [j^L, j^1] \dots, j^{\hat{l}_1}\}$ and $o^1 \in \{p^{\hat{l}_2}, \dots [p^L, p^1] \dots, p^{\hat{l}_1-1}\}$. If $m^* \geq \bar{m}$, then

$$(j^{\tilde{l}}, p^{\tilde{l}}, \dots [p^L, j^1] \dots, j^{l^*-1}, p^{l^*-1}, i^2, o^2, \dots, i^{\tilde{m}-1}, o^{\tilde{m}-1})$$

is a CIC of μ for $j^{\tilde{l}}$ at \hat{A} .

If $p^{\tilde{l}} \in \Omega_{j^{\tilde{l}}} \setminus A_{j^{\tilde{l}}}$, we have $j^{\tilde{l}} < \hat{j}^*$ and $p^{\tilde{l}-1} \in \Omega_{j^{\tilde{l}}} \setminus A_{j^{\tilde{l}}}$. If $o^1 \notin \{p^{\hat{l}_2}, \dots [p^L, p^1] \dots, p^{\hat{l}_1-1}\}$, then

$$(i^{\tilde{m}}, o^{\tilde{m}}, \dots, i^{\hat{m}_2-1}, o^{\hat{m}_2-1}, j^{\hat{l}_2}, p^{\hat{l}_2}, \dots [p^L, j^1] \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1})$$

is a CIC of μ for $i^{\tilde{m}}$ at \hat{A} . If there exists $l^* \in \{\hat{l}_2 + 1, \dots [L, 1] \dots, \tilde{l}\}$ such that $p^{l^*-1} = o^1$, we have $p^{l^*} \in A_{j^*}$. If $m^* < \underline{m}$, then

$$(j^{l^*}, p^{l^*}, \dots [p^L, j^1] \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1}, i^{\tilde{m}}, o^{\tilde{m}}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at \hat{A} . Otherwise,

$$(i^2, o^2, \dots, i^{\hat{m}_2-1}, o^{\hat{m}_2-1}, j^{\hat{l}_2}, p^{\hat{l}_2}, \dots [p^L, j^1] \dots, j^{l^*-1}, p^{l^*-1})$$

is a CIC of μ for $j^{\tilde{l}}$ at \hat{A} .

Case 2: $\tilde{m} \in \{\underline{m} + 1, \dots, \bar{m} - 1\}$, $i^{\tilde{m}} < \hat{j}^*$, $o^{\tilde{m}-1} \in A_{i^{\tilde{m}}}$, and $o^{\tilde{m}} \in \Omega_{i^{\tilde{m}}} \setminus A_{i^{\tilde{m}}}$. We construct a CIC of $\mu + C$ for $i^{\tilde{m}}$ at A and thus, given that $i^{\tilde{m}} < \hat{j}^*$, contradict the assumption that \hat{j}^* is the highest priority agent for whom there exists a CIC of $\mu + C$ at A .

Note first that since $\tilde{m} > \underline{m}$ there exists a largest integer $\hat{m} \in \{\underline{m}, \dots, \tilde{m} - 1\}$ such that $i^{\hat{m}} \in I(\hat{C})$. Let \hat{l} be such that $j^{\hat{l}} = i^{\hat{m}}$ and note that, since C is a CIC and $\hat{m} \neq \tilde{m}$, $i^{\hat{m}} \neq i^{\tilde{m}}$. Next, let \tilde{l} be the first integer in the sequence $(\hat{l} + 1, \dots [L, 1] \dots, \hat{l} - 1)$ such that $j^{\tilde{l}} \in \{i^{\hat{m}+1}, \dots, i^{\bar{m}}\}$. Note that such an integer exists since $\tilde{m} < \bar{m}$ and $j^{\hat{l}} \in \{i^{\underline{m}}, \dots, i^{\tilde{m}-1}\}$. Let \tilde{m} be such that $i^{\tilde{m}} = j^{\tilde{l}}$.

If $i^{\tilde{m}} \notin \{j^{\hat{l}}, \dots [j^L, j^1] \dots, j^{\tilde{l}}\}$, then

$$(i^{\tilde{m}}, o^{\tilde{m}-1}, \dots, i^{\hat{m}+1}, o^{\hat{m}}, j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1}, i^{\tilde{m}}, o^{\tilde{m}-1}, \dots, i^{\hat{m}+1}, o^{\hat{m}}).$$

is a CIC of $\mu + C$ for $i^{\tilde{m}}$ at A .

For the remainder of the proof in Case 2 assume that $i^{\tilde{m}} \in \{j^{\hat{l}}, \dots [j^L, j^1] \dots, j^{\tilde{l}}\}$.

If $p^{\check{l}-1} \in \Omega_{j^{\check{l}}} \setminus A_{j^{\check{l}}}$, then analogs of our earlier arguments show

$$(i^{\check{m}}, o^{\check{m}-1}, \dots, i^{\check{m}+1}, o^{\check{m}}, j^{\check{l}}, p^{\check{l}}, \dots [p^L, j^1] \dots, j^{\check{l}-1}, p^{\check{l}-1})$$

is a CIC of $\mu + C$ for $i^{\check{m}}$ at A .

If $p^{\check{l}-1} \in A_{j^{\check{l}}}$, then since $i^{\check{m}} < \hat{j}^*$ and since \hat{C} is a CIC of $\mu + C$ for \hat{j}^* at A , we have $p^{\check{l}} \in A_{j^{\check{l}}}$. Analogs of our earlier arguments show that

$$(j^{\check{l}}, p^{\check{l}}, \dots [p^L, j^1] \dots, j^{\check{l}-1}, p^{\check{l}-1}, i^{\check{m}}, o^{\check{m}-1}, \dots, i^{\check{m}+1}, o^{\check{m}})$$

is a CIC of $\mu + C$ for $j^{\check{l}} = i^{\check{m}}$ at A .

Case 3: $\check{m} = \underline{m}$, $i^{\check{m}} \leq \hat{j}^*$, $o^{\check{m}-1} \in \Omega_{i^{\check{m}}} \setminus A_{i^{\check{m}}}$, $o^{\check{m}} \in A_{i^{\check{m}}}$, and $p^{\check{l}-1} \in \Omega_{i^{\check{m}}} \setminus A_{i^{\check{m}}}$. Note first that the premise of this case implies $p^{\check{l}-1} \notin O(C)$: Otherwise, since \hat{C} is a CIC of $\mu + C$, we have $p^{\check{l}-1} = o^{\check{m}}$, which is incompatible with $o^{\check{m}} \in A_{i^{\check{m}}}$ and $p^{\check{l}-1} \in \Omega_{i^{\check{m}}} \setminus A_{i^{\check{m}}}$. Let \hat{l} be the last integer in $(\check{l} + 1, \dots [L, 1] \dots, \check{l} - 1)$ such that $j^{\hat{l}} \in I(C) \setminus \{j^*\}$ and let \hat{m} be such that $i^{\hat{m}} = j^{\hat{l}}$. If $\{p^{\hat{l}}, \dots [p^L, p^1] \dots, p^{\hat{l}-1}\} \cap O(C) = \emptyset$, then similar arguments to those employed in Case 1 show that

$$(i^{\hat{m}}, o^{\hat{m}}, \dots, i^{\hat{m}-1}, o^{\hat{m}-1}, j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1})$$

is a CIC of μ for $i^{\hat{m}}$ at \hat{A} . If $\{p^{\hat{l}}, \dots [p^L, p^1] \dots, p^{\hat{l}-1}\} \cap O(C) \neq \emptyset$, the definition of \hat{l} and our earlier finding that $p^{\check{l}-1} \notin O(C)$ jointly imply $\{p^{\hat{l}}, \dots [p^L, p^1] \dots, p^{\hat{l}-1}\} \cap O(C) = \{o^1\}$. Let l^* be such that $p^{l^*-1} = o^1$ and note that $j^{l^*} = j^*$. Furthermore, since \hat{C} is a CIC for $\hat{j}^* > j^*$ in the case we consider here, $p^{l^*} \in A_{j^*}$ as well. Next, note first that

$$(i^2, o^2, \dots, i^{\hat{m}-1}, o^{\hat{m}-1}, j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{l^*-1}, p^{l^*-1})$$

is a CIC of μ for $i^{\hat{m}}$ at \hat{A} unless $m^* \leq \hat{m} - 1$. Since $\underline{m} = \check{m} < \hat{m} \leq \overline{m}$, Claim 7 implies $m^* < \check{m}$. But then,

$$(j^{l^*}, p^{l^*}, \dots [p^L, j^1] \dots, j^{\check{l}-1}, p^{\check{l}-1}, i^{\check{m}}, o^{\check{m}}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at \hat{A} .

Case 4: $\check{m} = \overline{m}$, $i^{\check{m}} \leq \hat{j}^*$, $o^{\check{m}-1} \in \Omega_{i^{\check{m}}} \setminus A_{i^{\check{m}}}$, $o^{\check{m}} \in A_{i^{\check{m}}}$, and $p^{\check{l}} \in A_{i^{\check{m}}}$. Let \hat{l} be the first integer in $(\check{l} + 1, \dots [L, 1] \dots, \check{l} - 1)$ such that $j^{\hat{l}} \in I(C) \setminus \{j^*\}$ and let \hat{m} be such that $i^{\hat{m}} = j^{\hat{l}}$. If $\{p^{\hat{l}}, \dots [p^L, p^1] \dots, p^{\hat{l}-1}\} \cap O(C) = \emptyset$, then, given that Claim 7 implies

$m^* \notin \{\hat{m}, \dots, \check{m} - 1\}$,

$$(j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1}, i^{\hat{m}}, o^{\hat{m}}, \dots, i^{\check{m}-1}, o^{\check{m}-1})$$

is a CIC of μ for $j^{\hat{l}} = i^{\hat{m}}$ at \hat{A} . If $p^{\hat{l}-1} \in O(C)$ and $\{p^{\hat{l}}, \dots [p^L, p^1] \dots, p^{\hat{l}-2}\} \cap O(C) = \emptyset$, we have $p^{\hat{l}-1} = o^{\hat{m}}$ and

$$(j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1}, i^{\hat{m}+1}, o^{\hat{m}+1}, \dots, i^{\check{m}-1}, o^{\check{m}-1})$$

is a CIC of μ for $j^{\hat{l}} = i^{\hat{m}}$ at \hat{A} . By the definition of \hat{l} , the only remaining possibility is that there exists a $l^* \in \{\hat{l} + 1, \dots [L, 1] \dots, \hat{l}\}$ such that $p^{l^*-1} = o^1$. Since \hat{C} is a CIC for $\hat{j}^* > j^*$, we have $p^{l^*} \in A_{j^*}$. Next, note first that

$$(i^2, o^2, \dots, i^{\check{m}-1}, o^{\check{m}-1}, j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{l^*-1}, p^{l^*-1})$$

is a CIC of μ for $i^{\hat{m}}$ at \hat{A} unless $m^* \leq \check{m} - 1$. Since $\underline{m} < \bar{m} = \check{m}$, Claim 7 implies $m^* < \underline{m}$. But then,

$$(j^{l^*}, p^{l^*}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1}, i^{\check{m}}, o^{\check{m}}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at \hat{A} if $p^{\hat{l}-1} \notin O(C)$ and

$$(j^{l^*}, p^{l^*}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1}, i^{\check{m}+1}, o^{\check{m}+1}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at \hat{A} if $p^{\hat{l}-1} \in O(C)$.

Case 5: $\check{m} \leq \underline{m}$, $i^{\check{m}} < \hat{j}^*$, $o^{\check{m}-1} \in \Omega_{i^{\check{m}}} \setminus \mathbf{A}_{i^{\check{m}}}$, $o^{\check{m}} \in \mathbf{A}_{i^{\check{m}}}$, and $\{p^{l^*-1}, p^{l^*}\} \subseteq A_{j^*}$.

We argue first that $m^* \geq \bar{m}$. Assume the contrary and let \tilde{l} be the first integer in $(l^* + 1, \dots [L, 1] \dots, l^* - 1)$ such that $j^{\tilde{l}} \in I(C)$. Note that such an integer exists since $(I(C) \cap I(\hat{C})) \setminus \{j^*\} \neq \emptyset$. If $p^{\tilde{l}-1} \notin O(C)$, let \tilde{m} be such that $i^{\tilde{m}} = j^{\tilde{l}}$ and consider

$$\tilde{C} \equiv (j^{l^*}, p^{l^*}, \dots [j^L, p^1] \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1}, i^{\tilde{m}}, o^{\tilde{m}}, \dots, i^M, o^M).$$

Since $p^{l^*} \in A_{j^*}$, analogs of our earlier arguments show that \tilde{C} is a CIC of μ for $j^* = j^{\tilde{l}}$ at A . Since we assume that $m^* \not\geq \bar{m}$, Claim 7 implies $m^* < \underline{m}$. But then, we obtain that $m^* < \tilde{m}$ and \tilde{C} is a CIC of μ for j^* at \hat{A} , which contradicts our assumption that $\mu \in \mathcal{M}^{i-1}(\hat{A})$ given that $j^* < i$. If $p^{\tilde{l}-1} = o^{\tilde{m}-1}$ for some \tilde{m} , an analogous argument applies.

Now let \hat{l} be the last integer in $(l^* + 1, \dots [L, 1] \dots, l^* - 1)$ such that $j^{\hat{l}} \in I(C)$.

If $j^{\hat{l}} = i^{\check{m}}$, the premise of Case 5 implies $\check{m} = \underline{m}$. If $p^{\hat{l}-1} \in \Omega_{i^{\check{m}}} \setminus A_{i^{\check{m}}}$, then we can apply our findings from Case 3 above. If $p^{\hat{l}-1} \in A_{i^{\check{m}}}$, then, since \hat{C} is a CIC for \hat{j}^* , we have $i^{\check{m}} < \hat{j}^*$ and $p^{\hat{l}} \in A_{i^{\check{m}}}$ as well. If $p^{l^*-1} \neq o^1$, then $m^* \geq \bar{m}$ implies that

$$(j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{l^*-1}, p^{l^*-1}, i^1, o^1, \dots, i^{\check{m}-1}, o^{\check{m}-1}).$$

is a CIC of μ for $i^{\check{m}}$ at \hat{A} . If $p^{l^*-1} = o^1$, we obtain a similar contradiction using

$$(j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{l^*-1}, p^{l^*-1}, i^2, o^2, \dots, i^{\check{m}-1}, o^{\check{m}-1}).$$

For the remainder of the proof in Case 5, assume $j^{\hat{l}} \neq i^{\check{m}}$ and let \hat{m} be such that $i^{\hat{m}} = j^{\hat{l}}$. Since $\check{m} \in \{2, \dots, \underline{m}\}$, $\hat{m} > \check{m}$. If $p^{l^*-1} \neq o^1$, analogs of our earlier arguments show that

$$(i^1, o^1, \dots, i^{\hat{m}-1}, o^{\hat{m}-1}, j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots j^{l^*-1}, p^{l^*-1})$$

is a CIC of μ for $i^{\hat{m}}$ at \hat{A} . Again, if $p^{l^*-1} = o^1$, we obtain a similar contradiction using

$$(i^2, o^2, \dots, i^{\hat{m}-1}, o^{\hat{m}-1}, j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots j^{l^*-1}, p^{l^*-1}).$$

Case 6: $\check{m} \leq \underline{m}$, $i^{\check{m}} < \hat{j}^*$, $o^{\check{m}-1} \in A_{i^{\check{m}}}$, $o^{\check{m}} \in \Omega_{i^{\check{m}}} \setminus A_{i^{\check{m}}}$, and $\{p^{l^*-1}, p^{l^*}\} \subseteq A_{j^*}$. Let \hat{l} be the first integer in $(l^* + 1, \dots [L, 1] \dots, l^* - 1)$ such that $j^{\hat{l}} \in I(C)$.

If $j^{\hat{l}} = i^{\check{m}}$, the conditions of Case 6 imply that $\check{m} = \underline{m}$.

If $p^{\hat{l}-1} \in \Omega_{i^{\check{m}}} \setminus A_{i^{\check{m}}}$, then the definition of \hat{l} , our assumption about \check{m} , and $p^{l^*} \in A_{j^*}$ imply that

$$\tilde{C} \equiv (i^{\check{m}}, o^{\check{m}-1}, \dots, i^2, o^1, j^{l^*}, p^{l^*}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1}).$$

is a CIC of $\mu + C$ for $i^{\check{m}}$ at A . Since $i^{\check{m}} < \hat{j}^*$, we obtain a contradiction to our assumption that \hat{j}^* is the highest priority agent for whom there exists a CIC of $\mu + C$ at A .

If $p^{\hat{l}-1} \in A_{i^{\check{m}}}$, let \tilde{l} be the first integer in $(\hat{l} + 1, \dots [L, 1] \dots, \hat{l} - 1)$ such that $j^{\tilde{l}} \in I(C) \setminus \{j^*\}$ and let \tilde{m} be such that $i^{\tilde{m}} = j^{\tilde{l}}$. The definitions of \hat{l} and \tilde{l} together with our assumption that $\underline{m} \neq \bar{m}$ imply that $j^{\tilde{l}} \in I(C) \setminus \{i^{\check{m}}\}$. Hence, $\tilde{m} > \check{m}$ and

$$(j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1}, i^{\check{m}}, o^{\check{m}-1}, \dots, i^{\tilde{m}+1}, o^{\tilde{m}})$$

is a CIC of $\mu + C$ for $i^{\check{m}}$ at A .

For the remainder of the proof in Case 6, assume $j^{\hat{l}} \neq i^{\hat{m}}$ and let \hat{m} be such that $i^{\hat{m}} = j^{\hat{l}}$. Since $\check{m} \in \{2, \dots, \underline{m}\}$, $\check{m} < \hat{m}$. Analogs of our earlier arguments establish that

$$\tilde{C}' \equiv (i^{\hat{m}}, o^{\hat{m}-1}, \dots, i^2, o^1, j^{l^*}, p^{l^*}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1})$$

is a CIC of $\mu + C$ for $i^{\hat{m}}$ at A , which again contradicts our assumption that \hat{j}^* is the highest priority agent for whom there exists a CIC of $\mu + C$ at A .

Case 7: $\check{m} = \underline{m}$, $i^{\check{m}} < \hat{j}^*$, $o^{\check{m}-1} \in A_{i^{\check{m}}}$, $o^{\check{m}} \in \Omega_{i^{\check{m}}} \setminus A_{i^{\check{m}}}$, and $\{p^{\check{l}-1}, p^{\check{l}}\} \subseteq A_{i^{\check{m}}}$. By Case 6, we can assume that, if there exists l^* such that $j^{l^*} = j^*$, then $\{p^{l^*-1}, p^{l^*}\} \subseteq \Omega_{j^*} \setminus A_{j^*}$.

Now let \hat{l} be the first integer in $(\check{l} + 1, \dots [L, 1] \dots, \check{l} - 1)$ such that $j^{\hat{l}} \in I(C)$. If $j^{\hat{l}} \neq j^*$, let \hat{m} be such that $i^{\hat{m}} = j^{\hat{l}}$ and note that

$$(j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1}, i^{\hat{m}}, o^{\hat{m}-1}, \dots, i^{\check{m}+1}, o^{\check{m}})$$

is a CIC of $\mu + C$ for $i^{\hat{m}}$ at A . If $j^{\hat{l}} = j^*$, then

$$(j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{l^*-1}, p^{l^*-1}, j^{l^*}, o^M, i^M, o^{M-1}, \dots, i^{\check{m}+1}, o^{\check{m}})$$

is a CIC of $\mu + C$ for $i^{\hat{m}}$ at A .

Case 8: $\check{m} \geq \overline{m}$, $i^{\check{m}} < \hat{j}^*$, $o^{\check{m}-1} \in \Omega_{i^{\check{m}}} \setminus A_{i^{\check{m}}}$, $o^{\check{m}} \in A_{i^{\check{m}}}$, and $\{p^{l^*-1}, p^{l^*}\} \subseteq \Omega_{j^*} \setminus A_{j^*}$.

We argue first that $m^* < \underline{m}$. Assume the contrary and let \tilde{l} be the last integer in $(l^* + 1, \dots [L, 1] \dots, l^* - 1)$ such that $j^{\tilde{l}} \in I(C)$. Note that such an integer exists since $(I(C) \cap I(\hat{C})) \setminus \{j^*\} \neq \emptyset$. Let \tilde{m} be such that $i^{\tilde{m}} = j^{\tilde{l}}$. If $p^{\tilde{l}-1} \notin O(C)$, consider

$$\tilde{C} \equiv (i^1, o^1, \dots, i^{\tilde{m}-1}, o^{\tilde{m}-1}, j^{\tilde{l}}, p^{\tilde{l}}, \dots [j^L, p^1] \dots, j^{l^*-1}, p^{l^*-1}).$$

Since $p^{l^*-1} \in \Omega_{j^*} \setminus A_{j^*}$, analogs of our earlier arguments show that \tilde{C} is a CIC of μ for $j^* = i^1$ at A . Since we assume that $m^* \geq \underline{m}$, Claim 7 implies that $m^* \geq \overline{m}$. But then, we obtain that $m^* \geq \tilde{m}$ and \tilde{C} is a CIC of μ for j^* at \hat{A} , which contradicts our assumption that $\mu \in \mathcal{M}^{i-1}(\hat{A})$ given that $j^* < i$. If $p^{\tilde{l}-1} \in O(C)$, then $p^{\tilde{l}-1} = o^{\tilde{m}-1}$ and the same argument applies.

Now let \hat{l} be the first integer in $(l^* + 1, \dots [L, 1] \dots, l^* - 1)$ such that $j^{\hat{l}} \in I(C)$.

If $j^{\hat{l}} = i^{\check{m}}$, the premise of Case 8 implies $\check{m} = \overline{m}$. If $p^{\hat{l}} \in A_{i^{\check{m}}}$, then we can apply our findings from Case 4 above. If $p^{\hat{l}} \in \Omega_{i^{\check{m}}} \setminus A_{i^{\check{m}}}$, then, since \tilde{C} is a CIC for \hat{j}^* , we have $i^{\check{m}} < \hat{j}^*$ and $p^{\hat{l}-1} \in \Omega_{i^{\check{m}}} \setminus A_{i^{\check{m}}}$ as well. If $p^{l^*} \neq o^M$ in the case we consider here, then

$m^* < \underline{m}$ implies that

$$(i^{\check{m}}, o^{\check{m}}, \dots, i^M, o^M, j^{l^*}, p^{l^*}, \dots, j^{\hat{l}-1}, p^{\hat{l}-1})$$

is a CIC of μ for $i^{\check{m}}$ at \hat{A} . If $p^{l^*} = o^M$, we obtain a similar contradiction using

$$(i^{\check{m}}, o^{\check{m}}, \dots, i^M, o^M, j^{l^*+1}, p^{l^*+1}, \dots, j^{\hat{l}-1}, p^{\hat{l}-1}).$$

For the remainder of the proof in Case 8, we assume that $j^{\hat{l}} \neq i^{\check{m}}$ and let \hat{m} be such that $i^{\hat{m}} = j^{\hat{l}}$.

Since $\check{m} \in \{\overline{m}, \dots, M\}$, we have $\hat{m} < \check{m}$. If $p^{\hat{l}-1} \notin O(C)$ and $p^{l^*} \neq o^M$, analogs of our earlier arguments show that

$$(i^{\hat{m}}, o^{\hat{m}}, \dots, i^M, o^M, j^{l^*}, p^{l^*}, \dots, j^{\hat{l}-1}, p^{\hat{l}-1})$$

is a CIC of μ for $i^{\hat{m}}$ at \hat{A} . If $p^{\hat{l}-1} \notin O(C)$ and $p^{l^*} = o^M$, we obtain a similar contradiction using

$$(i^{\hat{m}}, o^{\hat{m}}, \dots, i^M, o^M, j^{l^*+1}, p^{l^*+1}, \dots, j^{\hat{l}-1}, p^{\hat{l}-1}).$$

If $p^{\hat{l}-1} \in O(C)$ and $p^{l^*} \neq o^M$, analogs of our earlier arguments show that

$$(i^{\hat{m}+1}, o^{\hat{m}+1}, \dots, i^M, o^M, j^{l^*}, p^{l^*}, \dots, j^{\hat{l}-1}, p^{\hat{l}-1})$$

is a CIC of μ for $i^{\hat{m}}$ at \hat{A} . Finally, if $p^{\hat{l}-1} \in O(C)$ and $p^{l^*} = o^M$, we obtain a similar contradiction using

$$(i^{\hat{m}+1}, o^{\hat{m}+1}, \dots, i^M, o^M, j^{l^*+1}, p^{l^*+1}, \dots, j^{\hat{l}-1}, p^{\hat{l}-1}).$$

Case 9: $\check{m} \geq \overline{m}$, $i^{\check{m}} < \hat{j}^*$, $o^{\check{m}-1} \in A_{i^{\check{m}}}$, $o^{\check{m}} \in \Omega_{i^{\check{m}}} \setminus A_{i^{\check{m}}}$, and $\{p^{l^*-1}, p^{l^*}\} \subseteq \Omega_{j^*} \setminus A_{j^*}$.

Let \hat{l} be the last integer in $(l^* + 1, \dots, [L, 1] \dots, l^* - 1)$ such that $j^{\hat{l}} \in I(C)$.

If $j^{\hat{l}} = i^{\check{m}}$, the conditions of Case 9 imply that $\check{m} = \overline{m}$.

If $p^{\hat{l}} \in A_{i^{\check{m}}}$, then

$$(j^{\hat{l}}, p^{\hat{l}}, \dots, j^{l^*-1}, p^{l^*-1}, i^1, o^M, \dots, i^{\overline{m}+1}, o^{\overline{m}}).$$

\tilde{C} is a CIC of $\mu + C$ for $i^{\check{m}}$ at A . Since $i^{\check{m}} < \hat{j}^*$, we obtain a contradiction to our assumption that \hat{j}^* is the highest priority agent for whom there exists a CIC of $\mu + C$ at A .

If $p^{\hat{l}} \in \Omega_{i^{\tilde{m}}} \setminus A_{i^{\tilde{m}}}$ note first that $p^{\hat{l}-1} \in \Omega_{i^{\tilde{m}}} \setminus A_{i^{\tilde{m}}}$ as well. Let \tilde{l} be the last integer in $(\tilde{l} + 1, \dots [L, 1] \dots, \tilde{l} - 1)$ such that $j^{\tilde{l}} \in I(C) \setminus \{j^*\}$ and let \tilde{m} be such that $i^{\tilde{m}} = j^{\tilde{l}}$. The definitions of \hat{l} and \tilde{l} imply $j^{\hat{l}} \neq i^{\tilde{m}}$. Hence, we have that $\tilde{m} < \bar{m}$ and

$$(j^{\tilde{l}}, p^{\tilde{l}}, \dots [p^L, j^1] \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1}, i^{\tilde{m}}, o^{\tilde{m}-1}, \dots, i^{\tilde{m}+1}, o^{\tilde{m}})$$

is a CIC of $\mu + C$ for $i^{\tilde{m}}$ at A .

For the remainder of the proof in Case 9, assume that $j^{\hat{l}} \neq i^{\tilde{m}}$ and let \hat{m} be such that $i^{\hat{m}} = j^{\hat{l}}$. Since $\tilde{m} \in \{\bar{m}, \dots, M\}$, $\hat{m} < \tilde{m}$. Analogs of our earlier arguments establish that

$$\tilde{C}' \equiv (j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{l^*-1}, p^{l^*-1}, i^1, o^M, \dots, i^{\hat{m}+1}, o^{\hat{m}})$$

is a CIC of $\mu + C$ for $i^{\tilde{m}}$ at A , which contradicts our assumption that \hat{j}^* is the highest priority agent for whom there exists a CIC of $\mu + C$ at A .

Case 10: $\tilde{m} = \bar{m}$, $i^{\tilde{m}} < \hat{j}^*$, $o^{\tilde{m}-1} \in A_{i^{\tilde{m}}}$, $o^{\tilde{m}} \in \Omega_{i^{\tilde{m}}} \setminus A_{i^{\tilde{m}}}$, and $\{p^{\tilde{l}-1}, p^{\tilde{l}}\} \subseteq \Omega_{i^{\tilde{m}}} \setminus A_{i^{\tilde{m}}}$.

By Case 9, we can assume that, if there exists l^* such that $j^{l^*} = j^*$, then $\{p^{l^*-1}, p^{l^*}\} \subseteq A_{j^*}$.

Now let \hat{l} be the last integer in $(\tilde{l} + 1, \dots [L, 1] \dots, \tilde{l} - 1)$ such that $j^{\hat{l}} \in I(C)$. If $j^{\hat{l}} \neq j^*$, let \hat{m} be such that $i^{\hat{m}} = j^{\hat{l}}$ and note that

$$(i^{\tilde{m}}, o^{\tilde{m}-1}, \dots, i^{\hat{m}+1}, o^{\hat{m}}, j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1})$$

is a CIC of $\mu + C$ for $i^{\tilde{m}}$ at A . If $j^{\hat{l}} = j^*$, then

$$(i^{\tilde{m}}, o^{\tilde{m}-1}, \dots, i^2, o^1, j^{l^*}, p^{l^*}, \dots [p^L, j^1] \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1})$$

is a CIC of $\mu + C$ for $i^{\tilde{m}}$ at A .

Case 11: $\tilde{m} \in \{\underline{m}, \dots, \bar{m}\}$, $i^{\tilde{m}} < \hat{j}^*$, $\{o^{\tilde{m}-1}, o^{\tilde{m}}\} \subseteq A_{i^{\tilde{m}}}$, and $\{p^{\tilde{l}-1}, p^{\tilde{l}}\} \subseteq \Omega_{j^i} \setminus A_{j^i}$.

Let \hat{l} be the last integer in $(\tilde{l} + 1, \dots [L, 1] \dots, \tilde{l} - 1)$ such that $j^{\hat{l}} \in I(C) \setminus \{j^*\}$ and let \hat{m} be such that $i^{\hat{m}} = j^{\hat{l}}$.

If $\hat{m} < \tilde{m}$, then

$$(i^{\tilde{m}}, o^{\tilde{m}-1}, \dots, i^{\hat{m}+1}, o^{\hat{m}}, j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{\tilde{l}-1}, p^{\tilde{l}-1})$$

is a CIC of $\mu + C$ for $i^{\tilde{m}}$ at A .

If $\hat{m} > \check{m}$ and $\{p^{\hat{l}}, \dots [p^L, p^1] \dots, p^{\hat{l}-1}\} \cap O(C) = \emptyset$, then Claim 7 implies that

$$(i^{\hat{m}}, o^{\hat{m}}, \dots, i^{\hat{m}-1}, o^{\hat{m}-1}, j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1})$$

is a CIC of μ for $i^{\hat{m}}$ at \hat{A} . By the premise of this case and the definition of \hat{l} , the only remaining possibility is that there exists $l^* \in (\hat{l} + 1, \dots [L, 1] \dots, \check{l} - 1)$ such that $p^{l^*-1} = o^1$. But then,

$$(i^{\hat{m}}, o^{\hat{m}-1}, \dots, i^2, o^1, j^{l^*}, p^{l^*}, \dots [p^L, j^1] \dots, j^{\check{l}-1}, p^{\check{l}-1})$$

is a CIC of $\mu + C$ for $i^{\hat{m}}$ at A .

Case 12: $\check{m} \in \{\underline{m}, \dots, \overline{m}\}$, $i^{\check{m}} < \hat{j}^*$, $\{o^{\check{m}-1}, o^{\check{m}}\} \subseteq \Omega_{j^*} \setminus A_{i^{\check{m}}}$, and $\{p^{\check{l}-1}, p^{\check{l}}\} \subseteq A_i$.

Let \hat{l} be the first integer in $(\check{l} + 1, \dots [L, 1] \dots, \check{l} - 1)$ such that $j^{\hat{l}} \in I(C) \setminus \{j^*\}$ and let \hat{m} be such that $i^{\hat{m}} = j^{\hat{l}}$.

If $\hat{m} < \check{m}$ and $\{p^{\hat{l}}, \dots [p^L, p^1] \dots, p^{\hat{l}-1}\} \cap O(C) = \emptyset$, then Claim 7

$$(j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1}, i^{\hat{m}}, o^{\hat{m}}, \dots, i^{\hat{m}-1}, o^{\hat{m}-1})$$

is a CIC of μ for $j^{\hat{l}}$ at \hat{A} . If $p^{\hat{l}-1} \in O(C)$ and $\{p^{\hat{l}}, \dots [p^L, p^1] \dots, p^{\hat{l}-2}\} \cap O(C) = \emptyset$, we have $p^{\hat{l}-1} = o^{\hat{m}}$ and

$$(j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1}, i^{\hat{m}+1}, o^{\hat{m}+1}, \dots, i^{\hat{m}-1}, o^{\hat{m}-1})$$

is a CIC of μ for $j^{\hat{l}} = i^{\hat{m}}$ at \hat{A} . By the definition of \hat{l} , the only remaining possibility is that there exists a $l^* \in (\check{l} + 1, \dots [L, 1] \dots, \hat{l} - 1)$ such that $p^{l^*-1} = o^1$. If $m^* \geq \check{m}$, then

$$(j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{l^*-1}, p^{l^*-1}, i^2, o^2, \dots, i^{\hat{m}-1}, o^{\hat{m}-1})$$

is a CIC of μ for $j^{\hat{l}}$ at \hat{A} . Otherwise, Claim 7 implies $m^* < \underline{m}$. Note that $p^{l^*-1} = o^1$ implies $p^{l^*} \in A_{j^*}$. If $p^{\hat{l}-1} \notin O(C)$, then

$$(j^{l^*}, p^{l^*}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1}, i^{\hat{m}}, o^{\hat{m}}, \dots, i^M, o^M)$$

is a CIC of μ for j^* at \hat{A} . If $p^{\hat{l}-1} \in O(C)$, the same is true for

$$(j^{l^*}, p^{l^*}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1}, i^{\hat{m}+1}, o^{\hat{m}+1}, \dots, i^M, o^M).$$

If $\hat{m} > \check{m}$, then

$$(j^{\hat{l}}, p^{\hat{l}}, \dots [p^L, j^1] \dots, j^{\hat{l}-1}, p^{\hat{l}-1}, i^{\hat{m}}, o^{\hat{m}-1}, \dots, i^{\check{m}+1}, o^{\check{m}})$$

is a CIC of $\mu + C$ for $j^{\hat{l}}$ at A .

□

E Proofs from Section 5

Proof of Theorem 4.

To prove part 1 of the Theorem, we show by induction on t that the maximum flows of (V, E, q) correspond to $\mathcal{M}^t(A)$ after the t^{th} iteration of the for-loop on line 2 of Algorithm 1.

As a base case, we show that for (V, E, q) , where q is the initial capacity vector defined above, f is a maximum flow if and only if it corresponds to some matching $\mu \in \mathcal{M}^0(A)$. Take any matching $\mu \in \mathcal{M}^0(A)$. The definition of a matching and trichotomous preferences imply $|\mu(i)| = |\Omega_i|$ and $\mu(i) \subseteq \Omega_i \cup A_i$. Define a flow f by setting, for all $i \in I$, $f(S, i) = |\Omega_i|$, $f(i, i^A) = |\mu(i) \cap A_i|$, $f(i, i^U) = |\mu(i) \cap (\Omega_i \setminus A_i)|$, $f(i^A, o) = 1$ for all $o \in \mu(i) \cap A_i$, $f(i^U, o) = 1$ for all $o \in \mu(i) \cap (\Omega_i \setminus A_i)$, and, for all $o \in \cup_{i \in I} \mu(i)$, $f(o, T) = 1$. Clearly, f is a maximum flow in (V, E, q) . Conversely, take a maximum flow f in (V, E, q) and let μ be the matching that corresponds to f . Since all maximum flows have value $\sum_i |\Omega_i|$, we have $|\mu(i)| = |\Omega_i|$ for all $i \in I$. Since (V, E, q) contains no paths between i and an object in $O \setminus (\Omega_i \cup A_i)$, we have $\mu(i) \subseteq \Omega_i \cup A_i$. Hence, $\mu \in \mathcal{M}^0(A)$.

As an induction hypothesis, suppose that at the start of the t^{th} iteration of the loop, the maximum flows of (V, E, q) correspond to $\mathcal{M}^{t-1}(A)$. Let \hat{q} be the capacity at the end of the t^{th} iteration of the for-loop, f be a maximum flow in (V, E, \hat{q}) , and μ be the matching that corresponds to f . We show first that $\mu \in \mathcal{M}^t(A)$. By construction, $|\mu(t) \cap A_t| = |\Omega_t| - \hat{q}(i, i^U)$. Assume that there is a matching $\mu' \in \mathcal{M}^t(A)$ such that $|\mu'(t) \cap A_t| > |\mu(t) \cap A_t|$. Since the set of maximum flows of (V, E, q) corresponds to $\mathcal{M}^{t-1}(A)$ and since $\mu' \in \mathcal{M}^t(A)$, just as argued in the base case, μ' induces a maximum flow f' in (V, E, q) . Given that $|\mu'(t) \cap A_t| > |\mu(t) \cap A_t|$, f' is a maximum flow in (V, E, \tilde{q}) , where $\tilde{q}(t, t^U) = \hat{q}(t, t^U) - 1$ and $\tilde{q}(e) = \hat{q}(e)$ for all $e \in E \setminus \{(t, t^U)\}$. But then, the t^{th} iteration of the loop would have concluded with the capacity of (i, i^U) no higher than $\hat{q}(t, t^U) - 1$. This contradiction shows that $\mu \in \mathcal{M}^t(A)$. Conversely, take any matching $\hat{\mu} \in \mathcal{M}^t$ and let \hat{f} be the induced flow, defined as in the base case. By our previous arguments, we have $|\hat{\mu}(t) \cap A_t| = |\mu(t) \cap A_t|$. Since $|\hat{\mu}(t)| = |\mu(t)| = |\Omega_t|$, we obtain that $|\hat{\mu}(t) \setminus A_t| = |\mu(t) \setminus A_t| = \hat{q}(t, t^U)$. Given that $\mu \in \mathcal{M}^t \subseteq \mathcal{M}^{t-1}$ and that

$\hat{q}(e) = q(e)$ for all $e \in E \setminus \{(t, t^U)\}$, the induction hypothesis implies that \hat{f} is a maximum flow in (V, E, \hat{q}) .

We now turn to part 2 of the Theorem. Let $m = |O|$ and $n = |I|$. Then there are at most m edges out of i^A for each $i \in I$, so the Ford-Fulkerson algorithm computes MAXFLOW in $O(mn^2)$ time. We invoke MAXFLOW while minimizing the flow along (i, i^U) , so it suffices to call it $O(\log(k))$ times for each agent, where $k = \max_{i \in N} |\Omega_i|$. Thus, the complexity of IRP is $O(mn^3 \log(k))$. Fixing Ω_i for each $i \in N$ rather than taking it as a parameter, the complexity is $O(mn^3)$ since k is a constant. \square